

WHITTAKER PERIODS, MOTIVIC PERIODS, AND SPECIAL VALUES OF TENSOR PRODUCT L -FUNCTIONS

HARALD GROBNER AND MICHAEL HARRIS

ABSTRACT. Let \mathcal{K} be an imaginary quadratic field. Let Π and Π' be irreducible generic cohomological automorphic representations of $GL(n)/\mathcal{K}$ and $GL(n-1)/\mathcal{K}$, respectively. Each of them can be given two natural rational structures over number fields. One is defined by the rational structure on topological cohomology, the other is given in terms of the Whittaker model. The ratio between these rational structures is called a *Whittaker period*. An argument presented by Mahnkopf and Raghuram shows that, at least if Π is cuspidal and the weights of Π and Π' are in a standard relative position, the critical values of the Rankin-Selberg product $L(s, \Pi \times \Pi')$ are essentially algebraic multiples of the product of the Whittaker periods of Π and Π' . We show that, under certain regularity and polarization hypotheses, the Whittaker period of a cuspidal Π can be given a motivic interpretation, and can also be related to a critical value of the adjoint L -function of related automorphic representations of unitary groups. The resulting expressions for critical values of the Rankin-Selberg and adjoint L -functions are compatible with Deligne's conjecture.

CONTENTS

1. Introduction	1
2. Unitary groups and base change	6
3. Whittaker periods for the general linear group	12
4. Motives – An Interlude	25
5. Rational Eisenstein classes of abelian type	35
6. Period relations - The main results	39
References	49

1. INTRODUCTION

L -functions can be attached both to automorphic representations and to arithmetic objects such as Galois representations or motives, and one implication of the Langlands program is that L -functions of the second kind are examples of L -functions of the first kind. Very few results of arithmetic interest can be proved about the second kind of L -functions until they have been identified with automorphic L -functions. For example, there is an

2010 *Mathematics Subject Classification*. Primary: 11F67; Secondary: 11F41, 11F70, 11F75, 22E55.

Key words and phrases. cuspidal automorphic representation, Deligne's conjecture, special values, L -function, periods, rationality.

H.G. is supported in parts by the Austrian Science Fund (FWF) Erwin Schrödinger grant, J 3076-N13, and the FWF-project, project number P 25974-N25.

The research leading to these results has received funding from the European Research Council under the European Community's Seventh Framework Programme (FP7/2007-2013) / ERC Grant agreement nÂ° 290766 (AAMOT). This is a version of paper to appear in *J. Inst. Math. Jussieu*.

extraordinarily deep web of conjectures relating the values at integer points of arithmetic (motivic) L -functions to cohomological invariants of the corresponding geometric (motivic) objects. In practically all the instances¹ where these conjectures have been proved, automorphic methods have proved indispensable.

At the same time, there is a growing number of results on special values of automorphic L -functions that make no direct reference to arithmetic. Instead, the special values are written as algebraic multiples of complex invariants defined by means of representation theory. Examples relevant to the present paper include [36], [41], [40] and [17], where the complex invariants are defined for representations of cohomological type by reference to the uniqueness of Whittaker or Shalika models. The present paper continues this series, proving a version of the main results of [36], [41] and [40] for the Rankin-Selberg L -function of a pair (Π, Π') of a cuspidal automorphic representation Π of GL_n of cohomological type, and a cuspidal *or* abelian automorphic representation Π' of GL_{n-1} , also of cohomological type, where the general linear groups are over an imaginary quadratic field \mathcal{K} . Here, the notion “abelian automorphic” refers to a representation which is a tempered Eisenstein representation induced from a Borel subgroup of GL_{n-1} . In view of Raghuram’s recent preprint [40], which we received after writing a first version of this paper, the inclusion of such automorphic representations Π' is the new feature of this result.

Our cohomological result is stated below as Theorem 1.3. The main purpose of the present paper is to bring this purely automorphic result closer to the motivic expression for the critical values of the Rankin-Selberg L -function $L(s, \Pi \times \Pi')$ embodied in Deligne’s conjecture [9]. In what follows, we let c denote complex conjugation of \mathcal{K} . Assume for the moment that both Π and Π' are cuspidal. In order to identify $L(s, \Pi \times \Pi')$ with a motivic L -function, we make the following hypothesis (see Hypothesis 4.19 for details):

Hypothesis 1.1. Both Π and Π' can be obtained by base change from holomorphic discrete series representations of unitary groups of arbitrary signature.

Up to twisting by an integral or half-integral power of the global character $g \mapsto \|\det(g)\|$, we may assume that Π and Π' have unitary central character (so the central critical value of $L(s, \Pi \times \Pi')$ is at $s = \frac{1}{2}$). Then Hypothesis 1.1 implies the polarization condition

$$\Pi^\vee \simeq \Pi^c \quad \text{and} \quad (\Pi')^\vee \simeq (\Pi')^c.$$

Conversely, it has been known for some time ([33], see also [31]) that the polarization condition implies that, for any hermitian space W over \mathcal{K} of dimension n , Π descends to a stable L -packet $\{\pi\}_W$ of cuspidal automorphic representations of $G = U(W)$, of discrete series type at infinity, provided it descends locally at all finite places; a similar condition holds for Π' . The signature in Hypothesis 1.1 is the invariant of W at a fixed complex embedding of \mathcal{K} . In particular, Hypothesis 1.1 is automatic for Π (resp. Π') if n is odd (resp. even).

Under Hypothesis 1.1, the L -packet $\{\pi^\vee\}_W$ contains (at least) one member for which π_∞^\vee belongs to the holomorphic discrete series. Let the signature of W be $(j, n - j)$ and choose an arithmetically normalized vector ϕ_j in the chosen π^\vee . Define $P^{(j)}(\Pi)$ to be the square norm of ϕ_j for the (arithmetically normalized) Petersson inner product. The $P^{(j)}(\Pi)$ are the motivic invariants in terms of which we express the critical values of $L(s, \Pi \times \Pi')$.

¹Whether or not the analytic formulas for special values of the Riemann zeta function and Dirichlet L -series should be considered exceptions is beyond the scope of this introduction.

More precisely, the arithmetic normalization is defined with reference to the coherent cohomology of the Shimura variety attached to G , or rather to the similitude group $GU(W)$. It thus depends on a choice of extension of ϕ_j to a (holomorphic) modular form on the similitude group. We ignore this irritating but crucial detail in the introduction; it is addressed at length in Remark 4.33.

Here is our main theorem:

Theorem 1.2. *Let Π and Π' be cuspidal automorphic representations of $GL_n(\mathbb{A}_{\mathcal{K}})$ and $GL_{n-1}(\mathbb{A}_{\mathcal{K}})$ which are cohomological with respect to irreducible algebraic representations E_{μ} and E_{λ} , respectively, where μ (resp. λ) is the corresponding highest weight (i.e., an n -tuple (resp. $n-1$ -tuple) of integers), assumed to satisfy the equivalent conditions of Lem. 2.1 (i.e., hypothesis 2.3 for $\mathfrak{m} = 0$). We assume Π and Π' satisfy Hypothesis 1.1. Let $\mathbb{Q}(\Pi_f)$ and $\mathbb{Q}(\Pi'_f)$ be the fields of definition of the finite parts Π_f and Π'_f of Π and Π' , respectively. Furthermore, we assume that the parameters of Π and Π' are sufficiently regular:*

$$\mu_i - \mu_{i+1} \geq 2 \text{ for all } i \text{ and } \lambda_j - \lambda_{j+1} \geq 2 \text{ for all } j$$

Then there are finite extensions $E(\Pi)/\mathbb{Q}(\Pi_f)$ and $E(\Pi')/\mathbb{Q}(\Pi'_f)$ such that, for every critical point $s_0 = \frac{1}{2} + m$ of $L(s, \Pi \times \Pi')$ with $m \geq 0$, there is a non-zero constant $u(m, \Pi_{\infty}, \Pi'_{\infty})$, depending only on m, μ, λ (or equivalently on $m, \Pi_{\infty}, \Pi'_{\infty}$), such that

$$L\left(\frac{1}{2} + m, \Pi_f \times \Pi'_f\right) \sim_{\mathcal{K}E(\Pi)E(\Pi')} u(m, \Pi_{\infty}, \Pi'_{\infty}) \prod_{j=1}^{n-1} P^{(j)}(\Pi) \prod_{k=1}^{n-2} P^{(k)}(\Pi'),$$

where “ $\sim_{\mathcal{K}E(\Pi)E(\Pi')}$ ” means up to multiplication by an element in the composition of number fields $\mathcal{K}E(\Pi)E(\Pi')$. Equivalently, for every critical point $s_0 = n - 1 + m$ of $L(s, R(M(\Pi) \otimes M(\Pi')))$ with $m \geq 0$,

$$L(n - 1 + m, R(M(\Pi) \otimes M(\Pi'))) \sim_{\mathcal{K}E(\Pi)E(\Pi')} u(m, \Pi_{\infty}, \Pi'_{\infty}) \prod_{j=1}^{n-1} P^{(j)}(\Pi) \prod_{k=1}^{n-2} P^{(k)}(\Pi').$$

The fields $\mathbb{Q}(\Pi_f)$ and $\mathbb{Q}(\Pi'_f)$ are number fields by [6], Thm. 3.13. The role of the finite extensions $E(\Pi)$ and $E(\Pi')$, which can be taken to be cyclotomic fields, is explained below.

The proof involves three main ingredients. The first is the extension mentioned above of the results of Mahnkopf and Raghuram. In what follows, Π is cuspidal but Π' is either cuspidal or abelian automorphic. Since Π and Π' are cohomological, their finite parts Π_f and Π'_f have natural rational structures over the number fields $\mathbb{Q}(\Pi_f)$ and $\mathbb{Q}(\Pi'_f)$, respectively, defined by the rational structure on cohomology with coefficients in the local systems attached to E_{μ} and E_{λ} . But since they are tempered, they are both globally generic, and therefore Π_f and Π'_f have rational structures defined in terms of their Whittaker models. The two rational structures are necessarily proportional, because the representations are irreducible. The factors of proportionality, called *Whittaker periods*, belong to \mathbb{C}^{\times} . Details will be provided in subsequent sections. In what follows, $p(\Pi)$ and $p(\Pi')$ are the Whittaker periods. Here is a statement.

Theorem 1.3. *Let $\Pi = BC(\pi) \|\cdot\|^{\mathfrak{m}}$ be a cuspidal automorphic representation of $GL_n(\mathbb{A}_{\mathcal{K}})$, $\mathfrak{m} \in \mathbb{Z}$, and let $\Pi' = BC(\pi')$ be a cuspidal or abelian automorphic representation of $GL_{n-1}(\mathbb{A}_{\mathcal{K}})$ obtained by base change from unitary groups as in Sections 2.4 and 2.5. In particular, Π is cohomological with respect to E_{μ} and Π' is cohomological with respect to*

E_λ . We assume the parameters μ and λ satisfy hypothesis 2.3. Then for every critical point $s_0 = \frac{1}{2} + m$ of $L(s, \Pi \times \Pi')$ with $m \geq 0$, there is a non-zero constant $p(m, \Pi_\infty, \Pi'_\infty) \in \mathbb{C}^\times$, depending only on m, μ, λ (or equivalently $m, \Pi_\infty, \Pi'_\infty$), such that

(1) For all critical values $\frac{1}{2} + m \in \text{Crit}(\Pi \times \Pi')$ with $m \geq 0$ and every $\sigma \in \text{Aut}(\mathbb{C})$,

$$\sigma \left(\frac{L(\frac{1}{2} + m, \Pi_f \times \Pi'_f)}{p(\Pi)p(\Pi')p(m, \Pi_\infty, \Pi'_\infty)\mathcal{G}(\omega_{\Pi'_f,0})} \right) = \frac{L(\frac{1}{2} + m, \sigma\Pi_f \times \sigma\Pi'_f)}{p(\sigma\Pi)p(\sigma\Pi')p(m, \sigma\Pi_\infty, \sigma\Pi'_\infty)\mathcal{G}(\omega_{\sigma\Pi'_f,0})}.$$

(2)

$$L(\frac{1}{2} + m, \Pi_f \times \Pi'_f) \sim_{\mathbb{Q}(\Pi_f)\mathbb{Q}(\Pi'_f)} p(\Pi)p(\Pi')p(m, \Pi_\infty, \Pi'_\infty)\mathcal{G}(\omega_{\Pi'_f,0}),$$

where “ $\sim_{\mathbb{Q}(\Pi_f)\mathbb{Q}(\Pi'_f)}$ ” means up to multiplication by an element in the composition of number fields $\mathbb{Q}(\Pi_f)\mathbb{Q}(\Pi'_f)$.

The proof of this result follows the methods of [36], [41], [40]: the branching formula for the restriction of the representation E_μ to GL_{n-1} shows that the Rankin-Selberg integral for $GL_n \times GL_{n-1}$ can be interpreted as a cohomological cup product precisely for the critical values s_0 of the theorem. On the one hand, the cup product is rational on rational cohomology classes; on the other hand, the Rankin-Selberg integral is, up to an unspecified archimedean factor, a rational multiple of the Euler product on automorphic forms that are rational with respect to the Whittaker rational structure. Comparing these two rationality properties, we obtain the result; the constant $p(m, \Pi_\infty, \Pi'_\infty)$ accounts for the archimedean Rankin-Selberg integral.

The proof of Theorem 1.3 nevertheless involves two novelties. The first is specific to the method of this paper, and consists in the observation that the method applied in the earlier work remains valid when Π' is abelian automorphic; the reason for this choice will be made apparent momentarily. The second novelty is of more general import. The main contribution to the archimedean invariants $p(m, \Pi_\infty, \Pi'_\infty)$ is a Rankin-Selberg zeta integral involving explicit cohomological vectors. For nearly 40 years, results of this type, for GL_n with $n > 2$, were presented subject to a local non-vanishing hypothesis, which in our present situation comes down to the claim that the archimedean zeta integral does not equal zero for the choice of vectors forced upon us by the method. In 2011, however, Binyong Sun discovered an abstract representation-theoretic method that allowed him to prove that such integrals never vanish. In his companion paper [46] he applies this method to cohomological Rankin-Selberg zeta integrals for $GL_n(\mathbb{R}) \times GL_{n-1}(\mathbb{R})$ and for $GL_n(\mathbb{C}) \times GL_{n-1}(\mathbb{C})$. Without Sun’s breakthrough, the results of the present paper would still be conditioned on the non-vanishing hypothesis.

When Π' is the abelian automorphic representation attached to an $(n-1)$ -tuple of Hecke characters χ_i , the L -function $L(s, \Pi \times \Pi')$ factors as a product of $n-1$ L -functions of GL_n , twisted by the χ_i . When Π' is cohomological, these L -functions are of the type whose special values were considered in [21] (completed by [22]). The second main ingredient of the proof is the expression of these special values in terms of the $P^{(j)}(\Pi)$ introduced in 1.2, and in terms of period invariants attached to the χ_i . On the other hand, an abelian automorphic Π' is an Eisenstein representation. Shahidi’s calculation of Whittaker coefficients for Eisenstein series is the third main ingredient of the proof. It allows us to identify the Whittaker period $p(\Pi')$ with a motivic period attached to the collection of χ_i . A similar identification was already exploited in [36], for Eisenstein classes attached to other parabolic subgroups. In our case, the motivic expression for $p(\Pi')$ exactly cancels the terms in the expression calculated

in [21], yielding an expression (Thm. 6.7) for $p(\Pi)$ solely in terms of Petersson norms of holomorphic forms on Shimura varieties, multiplied by purely archimedean invariants that have not yet been calculated explicitly:

Theorem 1.4. *Let Π be a cuspidal automorphic representation of $GL_n(\mathbb{A}_{\mathcal{K}})$, which can be obtained by base change from unitary groups of all signatures (i.e., more precisely, satisfies Hypothesis 4.19). Assume that Π is cohomological with respect to the irreducible, finite-dimensional, algebraic representation E_{μ} of highest weight $\mu_G = (\mu_1, \dots, \mu_n; -\mu_n, \dots, -\mu_1)$ and suppose that $\mu_i - \mu_{i+1} \geq 2$ for all i , so that there is a highest weight $\lambda_G = (\lambda; \lambda^{\vee})$ for G'_{∞} and an integer $m > 0$ such that $s_0 = \frac{1}{2} + m$ is a critical value of $L(s, \Pi \otimes \Pi')$ for any cuspidal automorphic representation Π' of $GL_{n-1}(\mathbb{A}_{\mathcal{K}})$, which is cohomological with respect to E_{λ} . Then there is a non-zero constant $Z(\Pi_{\infty})$, which depends only on the local representation Π_{∞} and such that*

$$p(\Pi) \sim_{\mathcal{K}E(\Pi)} Z(\Pi_{\infty}) \prod_{j=1}^{n-1} P^{(j)}(\Pi).$$

The constant $Z(\Pi_{\infty})$, like the expression $u(m, \Pi_{\infty}, \Pi'_{\infty})$ in Thm. 1.3, is well-defined up to a non-zero scalar in \mathcal{K} . It should be mentioned that the latter theorem (or equivalently Thm. 6.7) is conditional on a global non-vanishing hypothesis. Verifying this hypothesis for the central critical value appears to be extremely difficult in general, but fortunately can always be satisfied provided the infinitesimal character of Π is sufficiently regular.

Putting together Theorems 1.3 and 1.4 with the results of [21], we easily obtain the main theorem, with the constant $u(m, \Pi_{\infty}, \Pi'_{\infty}) = p(m, \Pi_{\infty}, \Pi'_{\infty}) \cdot Z(\Pi_{\infty}) \cdot Z(\Pi'_{\infty})$. The individual terms in the product depend on normalizations of archimedean Whittaker models, as do the Whittaker periods $p(\Pi)$ and $p(\Pi')$ themselves, but the product is independent of normalizations, up to \mathcal{K}^{\times} . Following the announcement of these results, Jie Lin has identified $u(m, \Pi_{\infty}, \Pi'_{\infty})$ as the explicit power of $2\pi i$ predicted by the automorphic version of Deligne's conjecture described in section 4.2; see Theorem 6.11 for a precise statement.

In the absence of the regularity hypothesis of 1.2, we can prove a rather different kind of result, using Wei Zhang's recent proof of a version of Neal Harris's Ichino-Ikeda conjecture for unitary groups. Suppose Π and Π' satisfy the hypotheses of [53]; in particular, Π and Π' are both supercuspidal at some prime of \mathcal{K} split over \mathbb{Q} . Then the conclusion of Thm. 1.3 holds, with $p(\Pi)$ and $p(\Pi')$ replaced by the values at $s = 1$ of the adjoint L -functions of π and π' respectively, and with slightly different archimedean factors (See Cor. 6.25).

Theorem 1.5. *Let Π and Π' be cuspidal automorphic representations of $GL_n(\mathbb{A}_{\mathcal{K}})$ and $GL_{n-1}(\mathbb{A}_{\mathcal{K}})$ which are cohomological with respect to E_{μ} and E_{λ} , respectively, assumed to satisfy the equivalent conditions of Lemma 2.1. We assume $\Pi = BC(\pi)$ and $\Pi' = BC(\pi')$ are obtained by base change from definite unitary groups and that the irreducible unitary cuspidal automorphic representations π and π' satisfy the hypotheses of [53] (more precisely, Hypothesis 6.12). Then for every critical point $s_0 = \frac{1}{2} + m$ of $L(s, \Pi \times \Pi')$ with $m \geq 0$, there are a non-zero complex constant $a(m, \Pi_{\infty}, \Pi'_{\infty})$, depending only on the archimedean components of Π and Π' , and an integer $a(n)$ only depending on n , such that,*

$$L\left(\frac{1}{2} + m, \Pi_f \times \Pi'_f\right) \sim_{\mathcal{K}\mathbb{Q}(\pi)\mathbb{Q}(\pi')} a(m, \Pi_{\infty}, \Pi'_{\infty}) \mathcal{G}(\varepsilon_{\mathcal{K},f})^{a(n)} L(1, \pi_f, Ad) L(1, \pi'_f, Ad),$$

where " $\sim_{\mathcal{K}\mathbb{Q}(\pi)\mathbb{Q}(\pi')}$ " means up to multiplication by an element in the composition of number fields $\mathcal{K}\mathbb{Q}(\pi)\mathbb{Q}(\pi')$. Equivalently, for every critical point $s_0 = n - 1 + m$ of $L(s, R(M(\Pi) \otimes$

$M(\Pi'))$ with $m \geq 0$,

$$L(n-1+m, R(M(\Pi) \otimes M(\Pi'))) \sim_{\mathcal{K}\mathbb{Q}(\pi)\mathbb{Q}(\pi')} a(m, \Pi_\infty, \Pi'_\infty) \mathcal{G}(\varepsilon_{\mathcal{K},f})^{a(n)} L(1, \pi_f, Ad) L(1, \pi'_f, Ad).$$

While Deligne's conjecture provides an expression of the critical values in terms of Deligne periods, which are defined in terms of motives, and whose status is therefore at least partially hypothetical, the Petersson norms are invariants of authentically arithmetic objects. The purpose of Section 4 is to provide plausible reasons to identify Deligne's periods in our situation with the expressions derived using [21]. It should be noted that [21] was written with applications to self-dual cohomological automorphic representations in mind, although many of the results are valid more generally. For this reason, the standing self-duality hypothesis of [21] is sometimes invoked tacitly, simplifying some statements but making it difficult for the reader (including the author of [21], at more than 15 years' distance) to determine which statements need to be modified to treat the general case. We have therefore attempted in Section 4, and especially in 4.4, to provide complete statements with precise parameters. The authors hope that definitive statements, over general CM fields, will be available in the near future.

For the convenience of the reader, we wrote 1.2 from the automorphic, as well as from the motivic point of view. It is shown in Optimistic Comparison 4.32 that the period that appears in Thm. 1.2, namely

$$\prod_{j=1}^{n-1} P^{(j)}(\Pi) \prod_{k=1}^{n-2} P^{(k)}(\Pi'),$$

is at least formally consistent with Deligne's conjecture, provided the Hodge types of the motives attached to Π and Π' satisfy the relations derived from Lemma 2.1. The article [24] computes Deligne periods for general Rankin-Selberg products and obtains different products of the same basic invariants $P^{(j)}(\Pi)$ and $P^{(k)}(\Pi')$ (under the identifications of Optimistic Comparison 4.32). The final section explains how one might hope to express the critical values of general Rankin-Selberg products, using normalized Rankin-Selberg integrals of Eisenstein cohomology classes.

The Whittaker period invariants $p(\Pi)$ are defined by comparing two rational structures on Π_f , one coming from the global realization of Π_f in the cohomology of the locally symmetric space S_n attached to GL_n , the other from the Whittaker model. It is well-known, however, that Π_f occurs in a range of cohomological degrees $b_n \leq i \leq c_n$, and each one gives rise to one (or more) period invariants. Our $p(\Pi)$ is attached to the lowest cohomological degree b_n ; the proof of Theorem 1.3 is based on the numerical coincidence that $b_n + b_{n-1} - 1$ equals the dimension of S_{n-1} . In [47], Eric Urban showed that, when $n = 2$, the Whittaker period attached to Π in the top cohomological degree c_2 is related to a non-critical special value of one of the two Asai L -functions attached to Π . It can be deduced from the arguments of [53] and the work of Jacquet, Rallis, and Flicker cited there that the phenomenon Urban discovered remains valid for general n ; this is the subject of our forthcoming article with E. Lapid [15].

We would like to thank Jie Lin for her endurance and great help in finding a vicious sign mistake in section 4, which troubled us while proof-reading the paper. We gratefully acknowledge the help of F. Shahidi and W. Zhang, for their answers to our questions about different steps in the proof of the main theorem. A. Raghuram helped us clear up an apparent

contradiction between our results and Deligne's conjecture. We are especially grateful to Binyong Sun, for proving the non-vanishing theorem that makes our results unconditional and that will inevitably make the whole field of special values considerably more attractive.

2. UNITARY GROUPS AND BASE CHANGE

2.1. Some standing assumptions and notation.

2.1.1. *Number fields.* Let \mathcal{K}/\mathbb{Q} denote an imaginary quadratic extension of \mathbb{Q} . We write $S(\mathcal{K})$ for the set of places of \mathcal{K} and S for the set of places of \mathbb{Q} . The ring of adeles over \mathcal{K} (resp. over \mathbb{Q}) is denoted $\mathbb{A}_{\mathcal{K}}$ (resp. \mathbb{A}). The ring of integers of \mathcal{K} is denoted \mathcal{O} and we let $\mathfrak{D}_{\mathcal{K}}$ be the different of \mathcal{K}/\mathbb{Q} , i.e., $\mathfrak{D}_{\mathcal{K}}^{-1} = \{x \in \mathcal{K} : \text{Tr}_{\mathcal{K}/\mathbb{Q}}(x\mathcal{O}) \subset \mathbb{Z}\}$, where $\text{Tr}_{\mathcal{K}/\mathbb{Q}}$ is the trace map from \mathcal{K} to \mathbb{Q} . The normalized absolute value of $\mathbb{A}_{\mathcal{K}}$ is denoted $\|\cdot\|$. We extend the Hecke character $\varepsilon_{\mathcal{K}} : \mathbb{Q}^{\times} \backslash \mathbb{A}^{\times} \rightarrow \mathbb{C}^{\times}$, associated to the extension \mathcal{K}/\mathbb{Q} via class field theory, to a Hecke character $\eta : \mathcal{K}^{\times} \backslash \mathbb{A}_{\mathcal{K}}^{\times} \rightarrow \mathbb{C}^{\times}$. It is easy to see that η is unitary and $\eta_{\infty}(z) = z^t \bar{z}^{-t}$ for a certain $t = t(\eta) \in \frac{1}{2} + \mathbb{Z}$.

For convenience, we will also fix, once and for all, a non-trivial, continuous, additive character $\psi : \mathcal{K} \backslash \mathbb{A}_{\mathcal{K}} \rightarrow \mathbb{C}^{\times}$. To this end, let $\psi_{\mathbb{Q}}$ be the additive character of $\mathbb{Q} \backslash \mathbb{A}$, as in Tate's thesis, namely, $\psi_{\mathbb{Q}}(x) = e^{2\pi i \Lambda(x)}$ with the $\Lambda = \sum_{p \leq \infty} \Lambda_p$, where $\Lambda_{\infty}(x) = -x$ for $x \in \mathbb{R}$ and $\Lambda_p(x_p)$, $x_p \in \mathbb{Q}_p$, is the finite, negative tail of x_p 's p -adic expansion. That is, if $x_p = \sum_{j \geq v} a_j p^j$, then $\Lambda_p(x_p) = \sum_{j=v}^{-1} a_j p^j$. If we write $\psi_{\mathbb{Q}} = \psi_{\mathbb{R}} \otimes \otimes_p \psi_{\mathbb{Q}_p}$, then $\psi_{\mathbb{R}}(x) = e^{-2\pi i x}$ and $\psi_{\mathbb{Q}_p}$ is trivial on \mathbb{Z}_p and non-trivial on $p^{-1}\mathbb{Z}_p$.

Now, we define $\psi := \psi_{\mathbb{Q}} \circ \text{Tr}_{\mathcal{K}/\mathbb{Q}}$. If $\psi = \otimes_w \psi_w$, then the local characters are determined analogously. In particular, if $\mathfrak{D}_{\mathcal{K}} = \prod_{\wp} \wp^{r_{\wp}}$, the product running over all prime ideals $\wp \subset \mathcal{O}$, then the conductor of the local character ψ_w is $\wp_w^{-r_{\wp}}$, i.e., ψ_w is trivial on $\wp_w^{-r_{\wp}}$ and non-trivial on $\wp_w^{-r_{\wp}-1}$. We let $S(\psi) = \{\wp \mid \mathfrak{D}_{\mathcal{K}}\}$.

2.1.2. *Hermitian spaces and unitary groups.* We let V be an n -dimensional Hermitian space over \mathcal{K} , relative to the non-trivial element in $\text{Gal}(\mathcal{K}/\mathbb{Q})$. Depending on the parity of n , we define

$$\gamma := \begin{cases} \mathbf{1} & \text{if } n \text{ is even} \\ \eta & \text{if } n \text{ is odd} \end{cases}$$

Hence, $\gamma_{\infty}(z) = z^c \bar{z}^{-c}$, where $c = 0$, if n is even and $c = t \in \frac{1}{2} + \mathbb{Z}$, if n is odd.

We let $V' \subset V$ be a subspace of codimension one, on which the restriction of the Hermitian form is non-degenerate so that $V = V' \oplus V'^{\perp}$. At infinity, we want V (and V') to be definite. We let $H := U(V)$ and $H' := U(V')$ be the corresponding unitary groups over \mathbb{Q} . In particular, $H(\mathbb{R}) \cong U(n)$, the compact unitary group of rank n and similar $H'(\mathbb{R}) \cong U(n-1)$.

Furthermore, we let $G := GL_n/\mathcal{K}$ and $G' := GL_{n-1}/\mathcal{K}$. We also define real Lie groups $K := GU(V)(\mathbb{R})$ (resp. $K' := GU(V')(\mathbb{R})$), which we view inside $G(\mathbb{C})$ (resp. $G'(\mathbb{C})$) by their natural embedding.

The Lie algebra of a real Lie group is denoted by the same but gothic, lower case letter.

2.1.3. *Finite-dimensional representations of real groups.* Let E_{μ}^{unt} (resp. E_{λ}^{unt}) be an irreducible, finite-dimensional, algebraic representation of $H(\mathbb{R})$ (resp. $H'(\mathbb{R})$) on a complex vector space. Having fixed a maximal \mathbb{Q} -split torus in $H(\mathbb{C}) \cong GL_n(\mathbb{C})$ (resp. $H'(\mathbb{C}) \cong GL_{n-1}(\mathbb{C})$) and an ordering on the set of its dominant algebraic characters, we may think of E_{μ}^{unt} (resp. E_{λ}^{unt}) as being given by its highest weight μ (resp. λ). We may arrange that

$\mu = (\mu_1, \dots, \mu_n)$, where $\mu_1 \geq \dots \geq \mu_n$, $\mu_i \in \mathbb{Z}$ for all $1 \leq i \leq n$ and similar $\lambda = (\lambda_1, \dots, \lambda_{n-1})$ with $\lambda_1 \geq \dots \geq \lambda_{n-1}$, $\lambda_i \in \mathbb{Z}$ for all $1 \leq i \leq n-1$.

Furthermore, we write $E_\mu^{\text{unt}} \otimes E_{\mu^\vee}^{\text{unt}}$ (resp. $E_\lambda^{\text{unt}} \otimes E_{\lambda^\vee}^{\text{unt}}$) for the irreducible, finite-dimensional, algebraic representation of the *real* Lie group $G(\mathbb{C})$ (resp. $G'(\mathbb{C})$) of highest weight (μ, μ^\vee) (resp. (λ, λ^\vee)). (Here the check “ \vee ” denotes taking the contragredient representation.) It is isomorphic to its complex conjugate contragredient.

Lemma 2.1 (See, e.g., [12], Thm. 8.1.1). *With the above notation, the following assertions are equivalent:*

- (1) $\mu_1 \geq -\lambda_{n-1} \geq \mu_2 \geq -\lambda_{n-2} \geq \dots \geq -\lambda_1 \geq \mu_n$
- (2) $\text{Hom}_{H(\mathbb{C})}(E_\mu^{\text{unt}} \otimes E_\lambda^{\text{unt}}, \mathbb{C}) \neq 0$

If any of these two conditions is satisfied, then also $\text{Hom}_{G'(\mathbb{C})}((E_\mu^{\text{unt}} \otimes E_{\mu^\vee}^{\text{unt}}) \otimes (E_\lambda^{\text{unt}} \otimes E_{\lambda^\vee}^{\text{unt}}), \mathbb{C}) \neq 0$

2.2. Base change. We let π (resp. π') be an irreducible, unitary cuspidal automorphic representation of $H(\mathbb{A})$ (resp. $H'(\mathbb{A})$). We shall always assume that π (resp. π') is cohomological with respect to E_μ^{unt} (resp. E_λ^{unt}). Since $H(\mathbb{R})$ and $H'(\mathbb{R})$ are both compact and connected, this simply means that $\pi_\infty \cong E_{\mu^\vee}^{\text{unt}}$ and $\pi'_\infty \cong E_{\lambda^\vee}^{\text{unt}}$. We let $S(\pi)$ (resp. $S(\pi')$) be the set of all finite places of \mathbb{Q} , where π (resp. π') ramifies, together with the infinite place of \mathbb{Q} . The following is essentially a theorem of Labesse:

Theorem 2.2 (Base Change). *There exist irreducible automorphic representations Π of $G(\mathbb{A}_K)$ and Π' of $G'(\mathbb{A}_K)$, which are of the form*

$$\Pi \cong \Pi_1 \boxplus \dots \boxplus \Pi_s \quad \text{and} \quad \Pi' \cong \Pi'_1 \boxplus \dots \boxplus \Pi'_r$$

where Π_j (resp. Π'_j) is a square-integrable automorphic representation of $GL_{s_j}(\mathbb{A}_K)$ (resp. $GL_{r_j}(\mathbb{A}_K)$), for all j , such that:

- For all $p \notin S(\pi)$, the representation $\Pi_p := \otimes_{w|p} \Pi_w$ is the local base change of π_p .
- The archimedean component Π_∞ of Π is cohomological with respect to $E_\mu^{\text{unt}} \otimes E_{\mu^\vee}^{\text{unt}}$, i.e.,
 $H^*(\mathfrak{g}, K, \Pi_\infty \otimes (E_\mu^{\text{unt}} \otimes E_{\mu^\vee}^{\text{unt}})) \neq 0$.

and analogously

- For all $p \notin S(\pi')$, the representation $\Pi'_p := \otimes_{w|p} \Pi'_w$ is the local base change of π'_p .
- The archimedean component Π'_∞ of Π' is cohomological with respect to $E_\lambda^{\text{unt}} \otimes E_{\lambda^\vee}^{\text{unt}}$, i.e.,
 $H^*(\mathfrak{g}', K', \Pi'_\infty \otimes (E_\lambda^{\text{unt}} \otimes E_{\lambda^\vee}^{\text{unt}})) \neq 0$.

If all the isobaric summands Π_j (resp. Π'_j) are cuspidal, then Π_∞ (resp. Π'_∞) is the local base change of π_∞ (resp. π'_∞).

Proof. This is essentially due to Labesse, [33] Cor. 5.3. Strictly speaking, the proof given in the aforementioned reference only works for totally real number fields F different from \mathbb{Q} . However, there is no doubt that the theorem holds in the above form. See [33], Remarque 5.2 and also [31]. \square

2.3. Descent to unitary groups and base change. When $BC(\pi)$ is cuspidal, it is expected, and has been proved in many cases (see for example [33]) that π occurs *with multiplicity one* in the discrete spectrum of H . Moreover, if H^b is any inner form of H then Π descends to an L -packet $\{\pi^b\}$ of discrete (cuspidal) automorphic representations of H^b whose

archimedean components lie in the discrete series; again it is expected that each member of the packet occurs with multiplicity one.

Suppose $H^b(\mathbb{R}) \cong U(r, s)$, the unitary group of signature (r, s) . Then any π^b on H^b whose base change is $BC(\pi)$, and whose archimedean component is a holomorphic (discrete series) representation, will be denoted $\pi_{r,s}$. Thus, $\pi = \pi_{n,0}$ or $\pi_{0,n}$. It doesn't matter whether H is positive or negative definite; but when $rs > 0$, the difference between $\pi_{r,s}$ and $\pi_{s,r}$ needs to be respected, because they define holomorphic automorphic forms on non-isomorphic Shimura varieties (attached to the similitude groups containing H^b). The normalizations relevant to these Shimura varieties can be found in [21].

To $\pi_{r,s}$ we associate a period invariant $P^{(s)}(BC(\pi)) \in \mathbb{C}^\times$, as in §4.2 of [24]. Roughly speaking, $P^{(s)}(BC(\pi))$ is the square of the Petersson norm of an arithmetically normalized holomorphic automorphic form in $\pi_{r,s}$. Precise definitions are in [21]; see also section 4.4 below, especially Hypothesis 4.19, as well as Remark 6.9.

2.4. The cuspidal automorphic representation Π . Recall the automorphic representation $BC(\pi)$ from Thm. 2.2. We make the additional assumption that it is cuspidal and take an arbitrary, but henceforth fixed, integer $\mathfrak{m} \in \mathbb{Z}$. We define

$$\Pi := BC(\pi) \|\cdot\|^{\mathfrak{m}},$$

suppressing its dependence on \mathfrak{m} . By assumption, Π is cuspidal automorphic and hence Π is automatically globally ψ^{-1} -generic. We denote by $W(\Pi_f)$ the corresponding Whittaker model of Π_f , again suppressing the dependence on the fixed additive character ψ^{-1} . By Thm. 2.2, Π_∞ is cohomological with respect to the finite-dimensional, algebraic representation

$$E_\mu := (E_\mu^{\text{unt}} \cdot \det^{-\mathfrak{m}}) \otimes (E_{\mu^\vee}^{\text{unt}} \cdot \det^{-\mathfrak{m}}) = E_{\mu-\mathfrak{m}}^{\text{unt}} \otimes E_{\mu^\vee-\mathfrak{m}}^{\text{unt}}$$

of $G(\mathbb{C})$ of highest weight $\mu_G := (\mu - \mathfrak{m}, \mu^\vee - \mathfrak{m})$, hence, we have

$$\Pi_\infty \cong \text{Ind}_{B(\mathbb{C})}^{G(\mathbb{C})} [z_1^{\ell_1+\mathfrak{m}} \bar{z}_1^{-\ell_1+\mathfrak{m}} \otimes \dots \otimes z_n^{\ell_n+\mathfrak{m}} \bar{z}_n^{-\ell_n+\mathfrak{m}}],$$

where $B = TN$ is the standard Borel subgroup of G , and

$$\ell_j = -\mu_{n-j+1} + \frac{n+1}{2} - j.$$

In analogy to the case of unitary groups, we let $S(\Pi)$ be the set of finite places of \mathcal{K} , where Π ramifies, together with the infinite place of \mathcal{K} .

2.5. The abelian automorphic representation Π' . Now recall the automorphic representations $BC(\pi')$ from Thm. 2.2. We set $\Pi' := BC(\pi')$ and make the following additional assumption: We suppose that $\Pi' \cong \Pi'_1 \boxplus \dots \boxplus \Pi'_r$ is abelian (and therefore not square-integrable), i.e.,

$$\Pi'_j = BC(\chi_j) \cdot \gamma \quad 1 \leq j \leq n-1,$$

where $BC(\chi_j)$ is the base change of an algebraic Hecke character $\chi_j : U(1)(\mathbb{Q}) \backslash U(1)(\mathbb{A}) \rightarrow \mathbb{C}^\times$. More explicitly, $BC(\chi_j)(g) = \chi_j(g/\bar{g})$ for all $g \in \mathbb{A}_{\mathcal{K}}^\times$, where \bar{g} denotes conjugation relative to the non-trivial element in $\text{Gal}(\mathcal{K}/\mathbb{Q})$. At the archimedean place of \mathbb{Q} (resp. \mathcal{K}) we obtain $\chi_{j,\infty}(e^{i\theta}) = e^{ik_j\theta}$, $e^{i\theta} \in U(1)$, (resp. $BC(\chi_j)_\infty(z) = z^{k_j} \bar{z}^{-k_j}$, $z \in \mathbb{C}^\times$) for $k_j \in \mathbb{Z}$. Therefore,

$$\Pi'_\infty \cong \text{Ind}_{B'(\mathbb{C})}^{G'(\mathbb{C})} [z_1^{k_1+c} \bar{z}_1^{-k_1-c} \otimes \dots \otimes z_{n-1}^{k_{n-1}+c} \bar{z}_{n-1}^{-k_{n-1}-c}],$$

where $B' = T'N'$ is the standard Borel subgroup of G' and so, since Π'_∞ is cohomological with respect to the finite-dimensional, algebraic representation

$$E_\lambda := E_\lambda^{\text{unt}} \otimes E_{\lambda^\vee}^{\text{unt}}$$

of $G'(\mathbb{C})$ of highest weight $\lambda_{G'} := (\lambda, \lambda^\vee)$, necessarily,

$$k_j = -\lambda_{n-j} + \frac{n}{2} - j - c.$$

Hence, we can further see that $k_1 > k_2 > \dots > k_{n-1}$. In particular, this implies, using Shahidi [44] Prop. 7.1.3, Thm. 3.5.12 and Rem. 3.5.14, that Π' is globally ψ -generic, i.e., the ψ -Whittaker coefficient of Π' defines a non-vanishing intertwining $\Pi' \hookrightarrow \text{Ind}_{N'(\mathbb{A})}^{G'(\mathbb{A})}[\psi]$ (un-normalized induction). As for Π , we denote by $W(\Pi'_f)$ the corresponding Whittaker model of Π'_f and let $S(\Pi')$ be the set of finite places of \mathcal{K} , where Π' ramifies, together with the infinite place of \mathcal{K} .

We will make the following assumption, valid throughout the paper:

Hypothesis 2.3. The highest weights μ and λ satisfy

$$\mu_1 \pm \mathfrak{m} \geq -\lambda_{n-1} \geq \mu_2 \pm \mathfrak{m} \geq -\lambda_{n-2} \geq \dots \geq -\lambda_1 \geq \mu_n \pm \mathfrak{m}$$

Lem. 2.1 then guarantees that $\text{Hom}_{G'(\mathbb{C})}(E_\mu \otimes E_\lambda, \mathbb{C}) \neq 0$, which will be important from Sect. 3.6 on.

2.6. An action of $\text{Aut}(\mathbb{C})$. For a moment, let $k \geq 1$ be any integer and let E be an irreducible, finite-dimensional, algebraic representation of GL_k/\mathcal{K} . As a representation of the real Lie group $GL_k(\mathbb{C})$, E factors as $E = E_\iota \otimes E_{\bar{\iota}}$, for a fixed embedding $\iota : \mathcal{K} \hookrightarrow \mathbb{C}$. For a given $\sigma \in \text{Aut}(\mathbb{C})$, we define the $GL_k(\mathbb{C})$ -representation

$${}^\sigma E := E_{\sigma \circ \iota} \otimes E_{\overline{\sigma \circ \iota}}.$$

Hence, ${}^\sigma E$ is identical to E , if $\sigma \circ \iota = \iota$ and it is the representation ${}^\sigma E = E_{\bar{\iota}} \otimes E_\iota$, otherwise, i.e., if $\sigma \circ \iota = \bar{\iota}$. Furthermore, for a representation of $GL_k(\mathbb{C})$, which is induced from the Borel subgroup $B_k = T_k N_k$ of GL_k , $\Delta_\infty = \text{Ind}_{B_k(\mathbb{C})}^{GL_k(\mathbb{C})}[\omega_1 \otimes \dots \otimes \omega_k]$, we let

$${}^\sigma \Delta_\infty := \text{Ind}_{B_k(\mathbb{C})}^{GL_k(\mathbb{C})}[{}^\sigma \omega_1 \otimes \dots \otimes {}^\sigma \omega_k].$$

Note that if Δ_∞ is cohomological with respect to E , then ${}^\sigma \Delta_\infty$ is cohomological with respect to ${}^\sigma E$.

If Δ_f (resp. Δ_w , $w \in S(\mathcal{K})_f$) is an irreducible representation of $GL_k(\mathbb{A}_f)$ (resp. $GL_k(\mathcal{K}_w)$), then we denote by ${}^\sigma \Delta_f$ (resp. ${}^\sigma \Delta_w$) the representation defined by Waldspurger in [48], I.1. Analogously, if δ_∞ is an irreducible representation of $H(\mathbb{R})$ or $H'(\mathbb{R})$, we let ${}^\sigma \delta_\infty := \delta_\infty$, since there is only the identity embedding of \mathbb{Q} into \mathbb{C} . For representations δ_f (resp. δ_p , $p \in S_f$) of $H(\mathbb{A}_f)$ or $H'(\mathbb{A}_f)$ (resp. $H(\mathbb{Q}_p)$ or $H'(\mathbb{Q}_p)$), we use again the definition of Waldspurger, [48], I.1.

If δ is any of the above representations, we let $\mathfrak{S}(\delta) := \{\sigma \in \text{Aut}(\mathbb{C}) \mid {}^\sigma \delta \cong \delta\}$ and define the rationality field of δ to be

$$\mathbb{Q}(\delta) := \{z \in \mathbb{C} \mid \sigma(z) = z \quad \forall \sigma \in \mathfrak{S}(\delta)\}.$$

See again Waldspurger, [48], I.1. We say that a representation δ is defined over a field F , if there is an F -subspace δ_0 of the representation space of δ , which is invariant under the group action and such that the natural map $\delta_0 \otimes_F \mathbb{C} \rightarrow \delta$ is an isomorphism.

Proposition 2.4. *For all $\sigma \in \text{Aut}(\mathbb{C})$, there is an isomorphism*

$$\sigma \Pi' \cong \boxplus_{j=1}^{n-1} \sigma BC(\chi_j) \sigma \left(\gamma \| \cdot \| \frac{n-2j}{2} \right) \| \cdot \| \frac{2j-n}{2}.$$

In other words, $\sigma \Pi'$ is again the isobaric automorphic sum of Hecke characters. Similarly, $\sigma \Pi$ is again a cuspidal automorphic representation.

Proof. The first assertion results from a straight-forward calculation, using the fact that since $BC(\chi_j)$ and $\gamma \| \cdot \| \frac{n-2j}{2}$ are both algebraic Hecke characters, so are $\sigma BC(\chi_j)$ and $\sigma \left(\gamma \| \cdot \| \frac{n-2j}{2} \right)$. The last assertion on Π can be shown as follows: By [6] Thm. 3.13, there is a cuspidal automorphic representation Ξ of $G(\mathbb{A}_{\mathcal{K}})$, cohomological with respect to σE_{μ} and such that $\Xi_f \cong \sigma \Pi_f$. As a cohomological cuspidal automorphic representation has an essentially tempered archimedean component, it follows from the classification of cohomological representations of $G(\mathbb{C})$ that necessarily $\Xi_{\infty} \cong \sigma \Pi_{\infty}$. \square

Proposition 2.5. *The finite part Π_f of $\Pi = BC(\pi) \| \cdot \| ^m$ (resp. Π'_f of $\Pi' = BC(\pi')$) is defined over its rationality field $\mathbb{Q}(\Pi_f)$ (resp. $\mathbb{Q}(\Pi'_f)$), which is an extension of $\mathbb{Q}(E_{\mu})$ (resp. of $\mathbb{Q}(E_{\lambda})$). This structure is unique up to homotheties. Both rationality fields are number fields. As a representation of $GL_k(\mathcal{K}) \subset GL_k(\mathbb{C})$, an irreducible, finite-dimensional, algebraic representation E of $GL_k(\mathbb{C})$ is defined over $\mathbb{Q}(E)$.*

Proof. The fields $\mathbb{Q}(\Pi_f)$ and $\mathbb{Q}(\Pi'_f)$ extend $\mathbb{Q}(E_{\mu})$, respectively $\mathbb{Q}(E_{\lambda})$, because of Strong Multiplicity One for isobaric automorphic representations. Here we use Prop. 2.4. Moreover, as a consequence of [6] Thm. 3.13, $\mathbb{Q}(\Pi_f)$ is a number field. (For a detailed proof, one may also have a look at [16], Thm. 8.1.) By Prop. 2.4

$$\bigcap_{j=1}^{n-1} \mathfrak{S} \left(BC(\chi_j)_f \gamma_f \| \cdot \|_f \frac{n-2j}{2} \right) \subseteq \mathfrak{S}(\Pi'_f),$$

so $\mathbb{Q}(\Pi'_f) \subseteq \prod_{j=1}^{n-1} \mathbb{Q} \left(BC(\chi_j)_f \gamma_f \| \cdot \|_f \frac{n-2j}{2} \right)$. As all $BC(\chi_j)$ and $\gamma \| \cdot \| \frac{n-2j}{2}$ are algebraic

Hecke characters, $\prod_{j=1}^{n-1} \mathbb{Q} \left(BC(\chi_j)_f \gamma_f \| \cdot \|_f \frac{n-2j}{2} \right)$ is a number field, whence so is $\mathbb{Q}(\Pi'_f)$.

The rest follows now from Clozel [6], Prop. 3.1 and p. 122. \square

Remark 2.6. The representation Π'_f is induced from the Hecke character $\beta := \Pi'_1 \otimes \cdots \otimes \Pi'_r$ of the (Levi quotient $T'(\mathbb{A}_f)$ of the) Borel subgroup $B'(\mathbb{A}_f)$, and can thus be identified with a space of functions on $G'(\mathbb{A}_f)$:

$$\Pi'_f = \{ \phi : G'(\mathbb{A}_f) \rightarrow \mathbb{C} \mid \phi(b'g') = \beta(b') \cdot \delta_{B'}^{\frac{1}{2}}(b') \cdot \phi(g'), \ b' \in B'(\mathbb{A}_f), \ g' \in G'(\mathbb{A}_f) \}.$$

Here $\delta_{B'}$ is the modulus character of $B'(\mathbb{A}_f)$. Define $\gamma^+(x) = \gamma(x) \cdot \|x\|^{\frac{1}{2}}$ if n is odd, $\gamma^+(x) = 1$ if n is even. For any n , $(\Pi'_j)^+ = BC(\chi_j) \cdot \gamma^+$ is an algebraic Hecke character, defined over the number field $\mathbb{Q}((\Pi'_j)^+)$. On the other hand, the j -th component of $\beta \cdot \delta_{B'}^{\frac{1}{2}}$ equals $BC(\chi_j) \cdot \gamma \| \cdot \| \frac{n-2j}{2}$ which is $(\Pi'_j)^+$ multiplied by an integral power of the norm character. Thus $\beta \cdot \delta_{B'}^{\frac{1}{2}}$ takes values in a number field, say $\mathbb{Q}(\beta, \gamma^+)$. It follows that the subspace

$$\Pi'_f[\mathbb{Q}(\beta, \gamma^+)] = \{ \phi : G'(\mathbb{A}_f) \rightarrow \mathbb{Q}(\beta, \gamma^+) \mid \phi(b'g') = \beta(b') \cdot \delta_{B'}^{\frac{1}{2}}(b') \cdot \phi(g'), \ b' \in B'(\mathbb{A}_f), \ g' \in G'(\mathbb{A}_f) \}$$

is a $G'(\mathbb{A}_f)$ -invariant $\mathbb{Q}(\beta, \gamma^+)$ -rational structure on Π'_f , and it is clear that the action of $\text{Aut}(\mathbb{C})$ on Π'_f can be read directly in terms of the actions on the values of functions.

Next, let $\mathbf{t}_{\sigma,k}$ be the diagonal matrix $\text{diag}(t_\sigma^{-(k-1)}, t_\sigma^{-(k-2)}, \dots, 1) \in GL_k(\mathbb{A}_f)$, as in [42], 3.2 or [36], 3.3. Here, t_σ is an element of $\widehat{\mathcal{O}}^*$, assigned to $\sigma \in \text{Aut}(\mathbb{C})$ by the map $\text{Aut}(\mathbb{C}) \rightarrow \text{Gal}(\mathbb{Q}(\mu_\infty)/\mathbb{Q}) \rightarrow \widehat{\mathbb{Z}}^* \hookrightarrow \widehat{\mathcal{O}}^*$. Let $\xi_f \in \text{Ind}_{N_k(\mathbb{A}_f)}^{GL_k(\mathbb{A}_f)}[\phi_f]$ (unnormalized induction), ϕ an additive, continuous character $\mathcal{K} \backslash \mathbb{A}_{\mathcal{K}} \rightarrow \mathbb{C}^\times$. Then, ${}^\sigma \xi_f(g_f) := \sigma(\xi(\mathbf{t}_{\sigma,k} g_f))$ is again an element in $\text{Ind}_{N_k(\mathbb{A}_f)}^{GL_k(\mathbb{A}_f)}[\phi_f]$. The analogous definition applies locally at a finite place $w \in S(\mathcal{K})_f$.

Proposition 2.7. *The map $\xi_f \mapsto {}^\sigma \xi_f$ defines a σ -linear $G(\mathbb{A}_f)$ -equivariant isomorphism from $W(\Pi_f)$ onto $W({}^\sigma \Pi_f)$ as well as a σ -linear $G'(\mathbb{A}_f)$ -equivariant isomorphism from $W(\Pi'_f)$ onto $W({}^\sigma \Pi'_f)$. For any finite extension F of the rationality field in question, we have an F -structure on $W(\Pi_f)$ and $W(\Pi'_f)$ by taking the $\text{Aut}(\mathbb{C}/F)$ -invariants.*

Proof. For the cuspidal representation Π , this is [42], Lem. 3.2. The same proof goes through for Π' , using Jacquet–Piatetski-Shapiro–Shalika’s theorem (5.1.(i)) in [28]. \square

2.7. Rational structures on cohomological Harish-Chandra modules. The universal enveloping algebra $\mathcal{U}(\mathfrak{g}')$ has a canonical \mathbb{Q} -rational (vector space-)structure determined by the \mathbb{Q} -reductive group $R_{\mathcal{K}/\mathbb{Q}}(G')$. Moreover, K' has a compatible \mathbb{Q} -rational structure, and determines a \mathbb{Q} -rational Cartan decomposition $\mathfrak{g}' = \mathfrak{k}' \oplus \mathfrak{p}'$. It thus makes sense to define the field of rationality of the $(\mathcal{U}(\mathfrak{g}'), K')$ -module Π'_∞ as the field $\mathbb{Q}(\Pi'_\infty)$ fixed by the action of the subgroup of $\text{Aut}(\mathbb{C})$ that fixes Π'_∞ up to isomorphism. Now Π'_∞ is the $\mathcal{U}(\mathfrak{g}')$ -module induced from a $\mathbb{Q}(E_\lambda)$ -rational character of the Lie algebra of the Borel subgroup $B'(\mathbb{C})$. In fact, it is given by non-normalized induction from an algebraic character of the torus $T'(\mathbb{C})$, by the arguments already seen in 2.6. It follows that the subspace of Π'_∞ isotypic for the representation E_λ of K' is defined over the number field $\mathbb{Q}(E_\lambda)$. Using the \mathbb{Q} -rational structure on \mathfrak{p}' , we find that the relative Lie algebra cohomology complex $C^\bullet(\mathfrak{g}', K', \Pi'_\infty \otimes E_\lambda) = \text{Hom}_{K'}(\Lambda^\bullet(\mathfrak{p}'), \Pi'_\infty \otimes E_\lambda)$ has a natural $\mathbb{Q}(E_\lambda)$ -rational (vector space-)structure. Hence, the same holds true for (\mathfrak{g}', K') -cohomology.

Although the cuspidal automorphic representation Π is not globally induced, it follows from the classification of cohomological representations that the archimedean component Π_∞ is again isomorphic to the non-normalized induction to $G(\mathbb{C})$ of an algebraic character of the maximal torus $T(\mathbb{C})$, see Sect. 2.4. We find again that $C^\bullet(\mathfrak{g}, K, \Pi_\infty \otimes E_\mu) = \text{Hom}_K(\Lambda^\bullet(\mathfrak{p}), \Pi_\infty \otimes E_\mu)$ – and so also the space of (\mathfrak{g}, K) -cohomology – has a natural $\mathbb{Q}(E_\mu)$ -rational (vector space-)structure.

In subsequent chapters we will be working with the Whittaker models $W(\Pi_\infty)$ and $W(\Pi'_\infty)$. It follows from the above discussion that their subspaces of K (resp. K')-finite vectors have rational models over the appropriate fields. Choices of complex isomorphisms $i_\infty : \Pi_\infty \xrightarrow{\sim} W(\Pi_\infty)$ and $i'_\infty : \Pi'_\infty \xrightarrow{\sim} W(\Pi'_\infty)$ identify the rational models of the two sides, up to complex factors of proportionality that depend on the choices. In the proof of Proposition 5.7, the factor of proportionality is denoted $\Omega(\Pi'_\infty)$; it is attached to the explicit choice of i'_∞ defined by the Whittaker integral. It is tempting to use the Whittaker integral to define i_∞ as well. This would provide natural normalizations for all the archimedean constants that appear in our final formulas. However, other normalizations – for example,

normalizations in terms of the archimedean local zeta integrals – may turn out to be more natural.

3. WHITTAKER PERIODS FOR THE GENERAL LINEAR GROUP

3.1. Automorphic cohomology of locally symmetric spaces. We define

$$S_n := G(\mathcal{K}) \backslash G(\mathbb{A}_{\mathcal{K}}) / K \quad \text{and} \quad S_{n-1} := G'(\mathcal{K}) \backslash G'(\mathbb{A}_{\mathcal{K}}) / K'$$

Moreover, we let

$$\tilde{S}_{n-1} := G'(\mathcal{K}) \backslash G'(\mathbb{A}_{\mathcal{K}}) / U(n-1)$$

Observe that $K \cong U(n)\mathbb{C}^\times = U(n)\mathbb{R}_+$ and similarly $K' \cong U(n-1)\mathbb{C}^\times = U(n-1)\mathbb{R}_+$, where \mathbb{R}_+ denotes the topological connected component of the identity of the split component of the center of $G(\mathbb{C})$ and $G'(\mathbb{C})$. In this way, the group $U(n-1)$ in the definition of \tilde{S}_{n-1} is K' without the contribution of the center. Consider the map $\iota : G' \hookrightarrow G$, realizing an element $g' \in G'$ as block diagonal matrix $\text{diag}(g', 1) \in G$. If K_f is an open compact subgroup of $G(\mathbb{A}_f)$, $K'_f := \iota^{-1}(K_f)$ is open compact in $G'(\mathbb{A}_f)$ and we define

$$S_n(K_f) := G(\mathcal{K}) \backslash G(\mathbb{A}_{\mathcal{K}}) / K K_f \quad \text{and} \quad S_{n-1}(K'_f) := G'(\mathcal{K}) \backslash G'(\mathbb{A}_{\mathcal{K}}) / K' K'_f$$

and $\tilde{S}_{n-1}(K'_f) := G'(\mathcal{K}) \backslash G'(\mathbb{A}_{\mathcal{K}}) / U(n-1) K'_f$. Our finite-dimensional modules E_μ and E_λ naturally define sheaves \mathcal{E}_μ on S_n and \mathcal{E}_λ on S_{n-1} , respectively, and we let

$$H^q(S_n, \mathcal{E}_\mu) \quad \text{and} \quad H^q(S_{n-1}, \mathcal{E}_\lambda)$$

be the corresponding cohomology spaces. They carry a $G(\mathbb{A}_f)$ -, resp. a $G'(\mathbb{A}_f)$ -module structure. With respect to this module-structure, there are isomorphisms

$$H^q(S_n, \mathcal{E}_\mu) \cong H^q(\mathfrak{g}, K, \mathcal{A}_{\mathcal{J}}(G) \otimes E_\mu)$$

and

$$H^q(S_{n-1}, \mathcal{E}_\lambda) \cong H^q(\mathfrak{g}', K', \mathcal{A}_{\mathcal{J}'}(G') \otimes E_\lambda).$$

This needs some explanation: First, the space $\mathcal{A}_{\mathcal{J}}(G)$ (resp. $\mathcal{A}_{\mathcal{J}'}(G')$) denotes the space of all automorphic forms of $G(\mathbb{A}_{\mathcal{K}})$ (resp. $G'(\mathbb{A}_{\mathcal{K}})$) which are annihilated by some power of \mathcal{J} (resp. \mathcal{J}'). Here, \mathcal{J} (resp. \mathcal{J}') is the ideal of the center of the universal enveloping algebra $\mathcal{U}(\mathfrak{g}_{\mathbb{C}})$ (resp. $\mathcal{U}(\mathfrak{g}'_{\mathbb{C}})$), which is annihilated by the contragredient representation of E_μ (resp. E_λ). Now, since $U(n)\mathbb{C}^\times = U(n)\mathbb{R}_+$ and $U(n-1)\mathbb{C}^\times = U(n-1)\mathbb{R}_+$, the assertion follows from Franke, [10], Thm. 18.

From this we obtain decompositions

$$H^q(S_n, \mathcal{E}_\mu) \cong H^q_{cusp}(S_n, \mathcal{E}_\mu) \oplus H^q_{Eis}(S_n, \mathcal{E}_\mu)$$

and

$$H^q(S_{n-1}, \mathcal{E}_\lambda) \cong H^q_{cusp}(S_{n-1}, \mathcal{E}_\lambda) \oplus H^q_{Eis}(S_{n-1}, \mathcal{E}_\lambda)$$

as follows: The space $\mathcal{A}_{\mathcal{J}}(G)$ may be decomposed along the associate classes $\{P\}$ of parabolic \mathcal{K} -subgroups $P = LN$ of G and the cuspidal supports φ_P , i.e., associate classes of irreducible cuspidal automorphic representations $\tau = \tilde{\tau} e^{(d\Lambda, H_P(\cdot))}$ of $L(\mathbb{A}_{\mathcal{K}})$, with $\tilde{\tau}$ a unitary cuspidal automorphic representation and $\Lambda : \mathbb{R}_+^{\text{rank}(P)} \rightarrow \mathbb{C}^\times$ a Lie group character of the split component $\mathbb{R}_+^{\text{rank}(P)}$ of $L(\mathbb{C})$. Indeed, if $\mathcal{A}_{\mathcal{J}, \{P\}}(G)$ denotes the space of all automorphic forms in $\mathcal{A}_{\mathcal{J}}(G)$, which are negligible along every parabolic subgroup $Q \notin \{P\}$, and, moreover, if $\mathcal{A}_{\mathcal{J}, \{P\}, \varphi_P}(G)$ denotes the subspace of $\mathcal{A}_{\mathcal{J}, \{P\}}(G)$, which is generated as a

$G(\mathbb{A})$ -module by all possible holomorphic values or residues of all Eisenstein series attached to $\tilde{\tau}$, evaluated at the point $d\Lambda$, together with all their derivatives, then

$$\mathcal{A}_{\mathcal{J}}(G) = \bigoplus_{\{P\}} \mathcal{A}_{\mathcal{J},\{P\}}(G) = \bigoplus_{\{P\},\varphi_P} \mathcal{A}_{\mathcal{J},\{P\},\varphi_P}(G).$$

For a detailed description of this decomposition, we refer the reader to the original paper, Franke-Schwermer [11], 1.1–1.4. If we set

$$\begin{aligned} H_{cusp}^q(S_n, \mathcal{E}_\mu) &:= H^q(\mathfrak{g}, K, \mathcal{A}_{\mathcal{J},\{G\}}(G) \otimes E_\mu) \\ H_{Eis}^q(S_n, \mathcal{E}_\mu) &:= \bigoplus_{\{P\} \neq \{G\}} H^q(\mathfrak{g}', K', \mathcal{A}_{\mathcal{J},\{P\}}(G) \otimes E_\mu) \end{aligned}$$

we obtain the above decomposition of cohomology for G . We remark that because $\mathcal{A}_{\mathcal{J},\{G\}}(G)$ consists precisely of all cuspidal automorphic forms in $\mathcal{A}_{\mathcal{J}}(G)$, $H_{cusp}^q(S_n, \mathcal{E}_\mu)$ is called the cuspidal cohomology of G (with respect to E_μ), and since $\mathcal{A}_{\mathcal{J},\{P\}}(G)$ is defined by means of Eisenstein series, supported in $\{P\}$, $H_{Eis}^q(S_n, \mathcal{E}_\mu)$ is called the Eisenstein cohomology of G (with respect to E_μ). Clearly, putting a prime everywhere, gives the analogous result for the cohomology of G' .

Let $H_c^q(S_n, \mathcal{E}_\mu)$ (resp. $H_c^q(\tilde{S}_{n-1}, \mathcal{E}_\lambda)$) be the cohomology with compact support. As cusp forms are rapidly decreasing, one has $H_{cusp}^q(S_n, \mathcal{E}_\mu) \subseteq H_c^q(S_n, \mathcal{E}_\mu)$. Put $b_k := \frac{k(k-1)}{2}$. Then b_k is the smallest degree in which a cohomological generic automorphic representation of $GL_k(\mathbb{A}_{\mathcal{K}})$ has non-zero cohomology. Note that this degree is independent of the given representation, as well as of the finite-dimensional coefficient module. Furthermore,

$$H^{b_n}(\mathfrak{g}, K, W(\Pi_\infty) \otimes E_\mu) \cong \mathbb{C}$$

and also

$$H^{b_{n-1}}(\mathfrak{g}', K', W(\Pi'_\infty) \otimes E_\lambda) \cong \mathbb{C}.$$

3.2. A diagram. Similar to the ideas in Mahnkopf [36], Raghuram [41], [40] and Grobner-Raghuram [17], we are going to consider the following diagram:

$$\begin{array}{ccc} H_c^{b_n}(S_n, \mathcal{E}_\mu) \times H^{b_{n-1}}(S_{n-1}, \mathcal{E}_\lambda) & \xrightarrow{\iota \times p} & H_c^{b_n}(\tilde{S}_{n-1}, \mathcal{E}_\mu) \times H^{b_{n-1}}(\tilde{S}_{n-1}, \mathcal{E}_\lambda) \\ \uparrow & & \downarrow \wedge \\ H_{cusp}^{b_n}(S_n, \mathcal{E}_\mu) \times H_{Eis}^{b_{n-1}}(S_{n-1}, \mathcal{E}_\lambda) & & H_c^{b_n+b_{n-1}}(\tilde{S}_{n-1}, \mathcal{E}_\mu \otimes \mathcal{E}_\lambda) \\ \uparrow \Psi = \Psi_\Pi^{\text{cusp}} \times \Psi_{\Pi'}^{\text{Eis}} & & \downarrow \mathcal{T}^* \\ H^{b_n}(\mathfrak{g}, K, \Pi \otimes E_\mu) \times H^{b_{n-1}}(\mathfrak{g}', K', \Pi' \otimes E_\lambda) & & H_c^{b_n+b_{n-1}}(\tilde{S}_{n-1}, \mathbb{C}) \\ \uparrow \Theta_0 = \Theta_0^{\text{cusp}} \times \Theta_0^{\text{Eis}} & & \downarrow f \\ W(\Pi_f) \times W(\Pi'_f) & \xrightarrow{\text{Dia}} & \mathbb{C} \end{array}$$

We are going to define the various maps, which appear in this diagram, in the next sections. Observe that “Dia” denotes the composition of all these maps.

3.3. The map Θ_0 . This section is in analogy with Mahnkopf [36], Raghuram-Shahidi, [42], Raghuram [41], [40] and Grobner-Raghuram [17].

As a first step, we choose and fix generators of the one-dimensional spaces $H^{b_n}(\mathfrak{g}, K, W(\Pi_\infty) \otimes E_\mu)$ and $H^{b_{n-1}}(\mathfrak{g}', K', W(\Pi'_\infty) \otimes E_\lambda)$. They are of the form

$$[\Pi_\infty] := \sum_{\underline{i}=(i_1, \dots, i_{b_n})} \sum_{\alpha=1}^{\dim E_\mu} X_{\underline{i}}^* \otimes \xi_{\Pi_\infty, \underline{i}, \alpha} \otimes e_\alpha,$$

and

$$[\Pi'_\infty] := \sum_{\underline{j}=(j_1, \dots, j_{b_{n-1}})} \sum_{\beta=1}^{\dim E_\lambda} X_{\underline{j}}'^* \otimes \xi_{\Pi'_\infty, \underline{j}, \beta} \otimes e'_\beta,$$

where the following data has been fixed:

- (1) An ordered basis $\{X_j\}$ of $\mathfrak{g}/\mathfrak{k}$, which fixes the dual-basis $\{X_j^*\}$ for $(\mathfrak{g}/\mathfrak{k})^*$. By our concrete choice of K , \mathfrak{k} is defined over \mathbb{Q} , whence we may assume that $\{X_j\}$ is a \mathbb{Q} -basis. For $\underline{i} = (i_1, \dots, i_{b_n})$, let $X_{\underline{i}}^* = X_{i_1}^* \wedge \dots \wedge X_{i_{b_n}}^* \in \bigwedge^{b_n} (\mathfrak{g}/\mathfrak{k})^*$.
- (2) Elements $e_1, \dots, e_{\dim E_\mu}$, which form a $\mathbb{Q}(E_\mu)$ -basis of E_μ .
- (3) To each \underline{i} and α , $\xi_{\Pi_\infty, \underline{i}, \alpha} \in W(\Pi_\infty)$.

and

- (1) An ordered basis $\{X'_j\}$ of $\mathfrak{g}'/\mathfrak{k}'$, which fixes the dual-basis $\{X'_j{}^*\}$ for $(\mathfrak{g}'/\mathfrak{k}')^*$. Similar to the case above, by our concrete choice of K' , \mathfrak{k}' is defined over \mathbb{Q} , whence we may assume that $\{X'_j\}$ is a \mathbb{Q} -basis. For $\underline{j} = (j_1, \dots, j_{b_{n-1}})$, let $X_{\underline{j}}'^* = X'_{j_1}{}^* \wedge \dots \wedge X'_{j_{b_{n-1}}}{}^* \in \bigwedge^{b_{n-1}} (\mathfrak{g}'/\mathfrak{k}')^*$.
- (2) Elements $e'_1, \dots, e'_{\dim E_\lambda}$, which form a $\mathbb{Q}(E_\lambda)$ -basis of E_λ .
- (3) To each \underline{j} and β , $\xi_{\Pi'_\infty, \underline{j}, \beta} \in W(\Pi'_\infty)$.

We may assume that the bases are compatible in the sense, that $\{X_j\}$ extends $\{X'_j\}$ (along the embedding $\mathfrak{g}'/\mathfrak{k}' \hookrightarrow \mathfrak{g}/\mathfrak{k}$ defined by ι) and that the non-compact part of the center of \mathfrak{g}' is spanned by X_r , $r = \dim_{\mathbb{R}} \tilde{S}_{n-1}$. The choice of the generators $[\Pi_\infty]$ and $[\Pi'_\infty]$ fixes generators $[\sigma\Pi_\infty]$ and $[\sigma\Pi'_\infty]$ of the one-dimensional spaces $H^{b_n}(\mathfrak{g}', K', W(\sigma\Pi_\infty) \otimes \sigma E_\mu)$ and $H^{b_{n-1}}(\mathfrak{g}', K', W(\sigma\Pi'_\infty) \otimes \sigma E_\lambda)$ for all $\sigma \in \text{Aut}(\mathbb{C})$ as well as isomorphisms Θ^{cusp} and Θ^{Eis} :

$$\begin{aligned} W(\Pi_f) &\xrightarrow{\sim} W(\Pi_f) \otimes H^{b_n}(\mathfrak{g}, K, W(\Pi_\infty) \otimes E_\mu) \\ &\xrightarrow{\sim} H^{b_n}(\mathfrak{g}, K, W(\Pi) \otimes E_\mu) \\ &\xrightarrow{\sim} H^{b_n}(\mathfrak{g}, K, \Pi \otimes E_\mu), \end{aligned}$$

and

$$\begin{aligned} W(\Pi'_f) &\xrightarrow{\sim} W(\Pi'_f) \otimes H^{b_{n-1}}(\mathfrak{g}', K', W(\Pi'_\infty) \otimes E_\lambda) \\ &\xrightarrow{\sim} H^{b_{n-1}}(\mathfrak{g}', K', W(\Pi') \otimes E_\lambda) \\ &\xrightarrow{\sim} H^{b_{n-1}}(\mathfrak{g}', K', \Pi' \otimes E_\lambda), \end{aligned}$$

where the last map in the respective diagram denotes the inverse of the corresponding Fourier coefficient. Next, recall from Prop. 2.5 that the $G(\mathbb{A}_f)$ -module $H^{b_n}(\mathfrak{g}, K, \Pi \otimes E_\mu)$ is defined over $\mathbb{Q}(\Pi_f)$ and that the $G'(\mathbb{A}_f)$ -module $H^{b_{n-1}}(\mathfrak{g}', K', \Pi' \otimes E_\lambda)$ is defined over $\mathbb{Q}(\Pi'_f)$. Both structures are unique up to multiplication by non-zero complex numbers. This leads us to the following

Proposition 3.1 (The Whittaker-periods). *There are non-zero complex numbers $p(\Pi) = p(\Pi_f, [\Pi_\infty])$ and $p(\Pi') = p(\Pi'_f, [\Pi'_\infty])$, such that the normalized maps*

$$(3.2) \quad \Theta_0^{\text{cusp}} := p(\Pi)^{-1} \cdot \Theta^{\text{cusp}} \quad \text{and} \quad \Theta_0^{\text{Eis}} := p(\Pi')^{-1} \cdot \Theta^{\text{Eis}}$$

are $\text{Aut}(\mathbb{C})$ -equivariant, i.e.,

$$\begin{array}{ccc} W(\Pi_f) & \xrightarrow{\Theta_0^{\text{cusp}}} & H^{b_n}(\mathfrak{g}, K, \Pi \otimes E_\mu) \\ \downarrow & & \downarrow \\ W(\sigma\Pi_f) & \xrightarrow{\Theta_0^{\text{cusp}}} & H^{b_n}(\mathfrak{g}, K, \sigma\Pi \otimes \sigma E_\mu) \end{array}$$

and

$$\begin{array}{ccc} W(\Pi'_f) & \xrightarrow{\Theta_0^{\text{Eis}}} & H^{b_{n-1}}(\mathfrak{g}', K', \Pi' \otimes E_\lambda) \\ \downarrow & & \downarrow \\ W(\sigma\Pi'_f) & \xrightarrow{\Theta_0^{\text{Eis}}} & H^{b_{n-1}}(\mathfrak{g}', K', \sigma\Pi' \otimes \sigma E_\lambda) \end{array}$$

commute. The complex number $p(\Pi)$ (resp. $p(\Pi')$) is well-defined only up to multiplication by non-zero elements of the number field $\mathbb{Q}(\Pi_f)$ (resp. $\mathbb{Q}(\Pi'_f)$).

Proof. For Π this is shown in [42], Definition/Proposition 3.3. In order to obtain the result for Π' , we observe that the proof of [42], Definition/Proposition 3.3, resp. [17] Definition/Proposition 4.2.1 goes over word for word, keeping in mind our Prop. 2.7 and Jacquet–Piatetski-Shapiro’s Thm. (5.1.(i)) in [28]. \square

Finally, we set

$$\Theta_0 := \Theta_0^{\text{cusp}} \times \Theta_0^{\text{Eis}}.$$

3.4. The map $\Psi = \Psi_\Pi^{\text{cusp}} \times \Psi_{\Pi'}^{\text{Eis}}$. It is well-known and follows from our résumé in section 3.1 together with Multiplicity One that

$$H^{b_n}(\mathfrak{g}, K, \Pi \otimes E_\mu) = H^{b_n}(\mathfrak{g}, K, \mathcal{A}_{\mathcal{J}, \{G\}, \varphi_\Pi} \otimes E_\mu),$$

where φ_Π is a singleton, represented by the cuspidal automorphic representation Π . Hence, there is a natural embedding

$$\Psi_\Pi^{\text{cusp}} : H^{b_n}(\mathfrak{g}, K, \Pi \otimes E_\mu) \hookrightarrow H_{\text{cusp}}^{b_n}(S_n, \mathcal{E}_\mu).$$

It is the purpose of this section to construct an embedding

$$\Psi_{\Pi'}^{\text{Eis}} : H^{b_{n-1}}(\mathfrak{g}', K', \Pi' \otimes E_\lambda) \hookrightarrow H_{\text{Eis}}^{b_{n-1}}(S_{n-1}, \mathcal{E}_\lambda),$$

as well. This is more delicate. First, as a short remark, let us point out that – in contrast to the case of cuspidal cohomology – this is also a question of degrees of cohomology: For a *non-trivial* $G'(\mathbb{A}_f)$ -morphism $H^{b_{n-1}}(\mathfrak{g}', K', \Pi' \otimes E_\lambda) \rightarrow H_{\text{Eis}}^{b_{n-1}}(S_{n-1}, \mathcal{E}_\lambda)$ to exist, it is necessary that

$$q_{\max} := \min_{\substack{P' = a \text{ max.} \\ \text{parabolic}/\mathcal{K}}} \left(\frac{1}{2} \dim_{\mathbb{R}} N_{P'}(\mathbb{C}) \right) \leq b_{n-1}.$$

This follows from Grobner [14], Thm. 18, since Π' is not square-integrable. Since for any maximal parabolic subgroup P' of G' , the unipotent radical $N_{P'}$ is of dimension $\dim_{\mathbb{C}} N_{P'}(\mathbb{C}) =$

$m_1 \cdot m_2$, for some non-trivial partition $m_1 + m_2 = n - 1$, we have without loss of generality $m_1 \leq \frac{n-1}{2}$ and $m_2 \leq n - 2$. Hence,

$$q_{\max} \leq \frac{1}{2} \dim_{\mathbb{R}} N_{P'}(\mathbb{C}) = m_1 \cdot m_2 \leq \frac{(n-1)(n-2)}{2} = b_{n-1},$$

and therefore the above condition is always satisfied. We now construct such a non-trivial map and show that is an injection.

Since b_{n-1} is the minimal degree, in which $\Pi'_{\infty} \cong \text{Ind}_{B'(\mathbb{C})}^{G'(\mathbb{C})}[z_1^{k_1+c} \bar{z}_1^{-k_1-c} \otimes \dots \otimes z_{n-1}^{k_{n-1}+c} \bar{z}_{n-1}^{-k_{n-1}-c}]$ has non-vanishing (\mathfrak{g}', K') -cohomology with respect to E_{λ} , there is an isomorphism of one-dimensional \mathbb{C} -vector-spaces:

$$\begin{aligned} & H^{b_{n-1}}(\mathfrak{g}', K', \text{Ind}_{B'(\mathbb{C})}^{G'(\mathbb{C})}[z_1^{k_1+c} \bar{z}_1^{-k_1-c} \otimes \dots \otimes z_{n-1}^{k_{n-1}+c} \bar{z}_{n-1}^{-k_{n-1}-c}] \otimes E_{\lambda}) \\ & \xrightarrow{\sim} H^{b_{n-1}}(\mathfrak{g}', K', \text{Ind}_{B'(\mathbb{C})}^{G'(\mathbb{C})}[z_1^{k_1+c} \bar{z}_1^{-k_1-c} \otimes \dots \otimes z_{n-1}^{k_{n-1}+c} \bar{z}_{n-1}^{-k_{n-1}-c} \otimes S(\check{\mathfrak{a}}_{B',\mathbb{C}}^{G'})] \otimes E_{\lambda}), \end{aligned}$$

where $S(\check{\mathfrak{a}}_{B',\mathbb{C}}^{G'})$ is the symmetric algebra of the orthogonal complement $\check{\mathfrak{a}}_{B',\mathbb{C}}^{G'}$ of $\check{\mathfrak{a}}_{G',\mathbb{C}} = X^*(G') \otimes_{\mathbb{Z}} \mathbb{C}$ in $\check{\mathfrak{a}}_{B',\mathbb{C}} = X^*(T') \otimes_{\mathbb{Z}} \mathbb{C}$. Hence, if we write $\tilde{\tau} := BC(\chi_1)\gamma \otimes \dots \otimes BC(\chi_{n-1})\gamma$, then

$$H^{b_{n-1}}(\mathfrak{g}', K', \Pi' \otimes E_{\lambda}) \cong H^{b_{n-1}}(\mathfrak{g}', K', \text{Ind}_{B'(\mathbb{A}_{\mathcal{K}})}^{G'(\mathbb{A}_{\mathcal{K}})}[\tilde{\tau} \otimes S(\check{\mathfrak{a}}_{B',\mathbb{C}}^{G'})] \otimes E_{\lambda}).$$

The algebra $\mathfrak{a}_{B',\mathbb{C}}^{G'}$ operates trivially on $\tilde{\tau}$. Hence, one may check that $(B', \tilde{\tau}, 0, 0)$ is one of the quadruples, constructed in Grobner [14], 3.3. Let $\varphi_{B'}$ be the associate class of unitary cuspidal automorphic representations of $T'(\mathbb{A}_{\mathcal{K}})$, represented by $\tilde{\tau}$, so the summand $\mathcal{A}_{\mathcal{J}',\{B'\},\varphi_{B'}}(G')$ of the space of automorphic forms $\mathcal{A}_{\mathcal{J}'}(G')$, cf. Sect. 3.1, is well-defined. Interpreting the elements of the symmetric algebra as differential operators $\frac{\partial^m}{\partial \Lambda^m}$, we obtain an intertwining operator

$$\begin{aligned} \text{Eis}_{\Pi'} : \text{Ind}_{B'(\mathbb{A}_{\mathcal{K}})}^{G'(\mathbb{A}_{\mathcal{K}})}[\tilde{\tau} \otimes S(\check{\mathfrak{a}}_{B',\mathbb{C}}^{G'})] &\rightarrow \mathcal{A}_{\mathcal{J}',\{B'\},\varphi_{B'}}(G') \\ f \otimes \frac{\partial^m}{\partial \Lambda^m} &\mapsto \frac{\partial^m}{\partial \Lambda^m} (E(f, \Lambda))|_{\Lambda=0}, \end{aligned}$$

where $E(f, \Lambda)$ is the Eisenstein series attached to a K' -finite section $f \in \text{Ind}_{B'(\mathbb{A}_{\mathcal{K}})}^{G'(\mathbb{A}_{\mathcal{K}})}[\tilde{\tau}]$ at $\Lambda \in \mathfrak{a}_{B',\mathbb{C}}^{G'}$. Observe that this intertwining operator is well-defined, since all Eisenstein series are holomorphic at $\Lambda = 0$.

Proposition 3.3. *The $G'(\mathbb{A}_f)$ -homomorphism*

$$\Psi_{\Pi'} : H^{b_{n-1}}(\mathfrak{g}', K', \text{Ind}_{B'(\mathbb{A}_{\mathcal{K}})}^{G'(\mathbb{A}_{\mathcal{K}})}[\tilde{\tau} \otimes S(\check{\mathfrak{a}}_{B',\mathbb{C}}^{G'})] \otimes E_{\lambda}) \rightarrow H^{b_{n-1}}(\mathfrak{g}', K', \mathcal{A}_{\mathcal{J}',\{B'\},\varphi_{B'}}(G') \otimes E_{\lambda})$$

induced from $\text{Eis}_{\Pi'}$ is an injection.

Proof. As all Eisenstein series $E(f, \Lambda)$ attached to a K' -finite section $f \in \text{Ind}_{B'(\mathbb{A}_{\mathcal{K}})}^{G'(\mathbb{A}_{\mathcal{K}})}[\tilde{\tau}]$ are holomorphic at $\Lambda = 0$, $\Psi_{\Pi'}$ is injective by Schwermer [45], Satz 4.11 or Li-Schwermer [34], Thm. 3.3. \square

As a consequence of our discussion in Sect. 3.1, we obtain an injection of $G'(\mathbb{A}_f)$ -modules

$$\Psi_{\Pi'}^{\text{Eis}} : H^{b_{n-1}}(\mathfrak{g}', K', \Pi' \otimes E_{\lambda}) = H^{b_{n-1}}(\mathfrak{g}', K', \Pi'_{\infty} \otimes E_{\lambda}) \otimes \Pi'_f \hookrightarrow H_{\text{Eis}}^{b_{n-1}}(S_{n-1}, \mathcal{E}_{\lambda}).$$

With respect to the $\mathbb{Q}(E_\lambda)$ -rational structure on relative Lie algebra cohomology defined in 2.7, this defines a $\mathbb{Q}(\Pi'_f)$ -rational injection

$$\Pi'_f = \text{Ind}_{B'(\mathbb{A}_f)}^{G'(\mathbb{A}_f)}[\tilde{\tau}_f] \hookrightarrow H_{\text{Eis}}^{b_{n-1}}(S_{n-1}, \mathcal{E}_\lambda)$$

where, for any field $L \supseteq \mathbb{Q}(\Pi'_f)$, the L -rational vectors on the left-hand side are just the L -valued functions on $G'(\mathbb{A}_f)$ that transform on the left under the character $\delta_{B',f}^{\frac{1}{2}} \cdot \tilde{\tau}_f$.

3.5. The map $\iota \times \phi$. Recall the map $\iota : G' \hookrightarrow G$, realizing an element $g' \in G'$ as block diagonal matrix $\text{diag}(g', 1) \in G$.

Lemma 3.4. *The map $\iota : \tilde{S}_{n-1}(K'_f) \rightarrow S_n(K_f)$ is proper.*

Proof. For the general linear group over \mathbb{Q} , this is stated in Mahnkopf [36], p. 615, however, without proof. For sake of completeness, we sketch an argument here. Let

$$S_{n-1,n}(K_f) := M(F) \backslash M(\mathbb{A}) / (K \cap M(\mathbb{C}))(K_f \cap M(\mathbb{A}_f)),$$

where $M \cong GL_{n-1} \times GL_1$, viewed as block diagonal matrices. The map ι factors as

$$\tilde{S}_{n-1}(K'_f) \xrightarrow{j} S_{n-1,n}(K_f) \xrightarrow{u} S_n(K_f).$$

Clearly, j is proper. Hence, it suffices to show that u is proper. This follows from [1], Lem. 2.7. \square

As a consequence, we obtain a map in cohomology with compact support:

$$\iota^q : H_c^q(S_n(K_f), \mathcal{E}_\mu) \rightarrow H_c^q(\tilde{S}_{n-1}(K'_f), \mathcal{E}_\mu).$$

Similar, the projection $p : \tilde{S}_{n-1}(K'_f) \rightarrow S_{n-1}(K'_f)$ induces a map

$$p^q : H^q(S_{n-1}(K'_f), \mathcal{E}_\lambda) \rightarrow H^q(\tilde{S}_{n-1}(K'_f), \mathcal{E}_\lambda).$$

In our diagram, we let $\iota \times p$ be the direct limit (over all open compact subgroups K_f of $G(\mathbb{A}_f)$) of the maps $\iota^{b_n} \times p^{b_{n-1}}$.

3.6. Critical points. Recall the definition of a point $s = \frac{1}{2} + m$, $m \in \mathbb{Z}$ being *critical* for $L(s, \Pi \times \Pi')$ from Deligne, [9] Prop.-Déf. 2.3. We let

$$\text{Crit}(\Pi \times \Pi') \subset \frac{1}{2} + \mathbb{Z}$$

be the set of critical points of $L(s, \Pi \times \Pi')$.

Lemma 3.5. *If $s = \frac{1}{2} \pm m$, $m \geq 0$, is a critical point of $L(s, \Pi \times \Pi')$, then $\text{Hom}_{H'(\mathbb{C})}(E_{\mu-m-m}^{\text{unt}} \otimes E_\lambda^{\text{unt}}, \mathbb{C}) \neq 0$ and $\text{Hom}_{H'(\mathbb{C})}(E_{\mu^v-m-m}^{\text{unt}} \otimes E_{\lambda^v}^{\text{unt}}, \mathbb{C}) \neq 0$. The set of critical points is $\text{Aut}(\mathbb{C})$ -invariant, i.e., $\text{Crit}(\Pi \times \Pi') = \text{Crit}(\sigma\Pi \times \sigma\Pi')$.*

Proof. A proof of these facts will be given in Sect. 4 in the motivic context. See in particular the proof of Lem. 4.7. \square

3.7. The map \mathcal{T}^* . Let $s = \frac{1}{2} + m \in \text{Crit}(\Pi \times \Pi')$, $m \geq 0$. Hence, by Lem. 3.5, we obtain a non-trivial map

$$\mathcal{T}^{(m)} \in \text{Hom}_{G'(\mathbb{C})}(E_{\mu-m} \otimes E_\lambda, \mathbb{C}),$$

which we will fix in a compatible way (i.e., $\mathcal{T}^{(m)}$ for $\Pi \otimes \Pi'$ shall be identical to $\mathcal{T}^{(0)}$ for $(\Pi \|\cdot\|^m) \otimes \Pi'$). Again by Lem. 3.5, it factors as $\mathcal{T}^{(m)} = \mathcal{T}_\iota^{(m)} \otimes \mathcal{T}_{\bar{\iota}}^{(m)}$, and if $\sigma \in \text{Aut}(\mathbb{C})$, then we define ${}^\sigma \mathcal{T}^{(m)}$ in the obvious way, i.e., ${}^\sigma \mathcal{T}^{(m)} = \mathcal{T}_{\sigma\circ\iota}^{(m)} \otimes \mathcal{T}_{\overline{\sigma\circ\iota}}^{(m)}$. Finally, if $s = \frac{1}{2} \in \text{Crit}(\Pi \times \Pi')$, we let $\mathcal{T} = \mathcal{T}^{(0)}$ and obtain a morphism in cohomology

$$\mathcal{T}^* : H_c^{b_n+b_{n-1}}(\tilde{S}_{n-1}, \mathcal{E}_\mu \otimes \mathcal{E}_\lambda) \rightarrow H_c^{b_n+b_{n-1}}(\tilde{S}_{n-1}, \mathbb{C}),$$

as in our diagram.

3.8. Poincaré duality. It remains to define the last map, denoted “ \int ” in our diagram. Therefore, observe that

$$b_n + b_{n-1} = \dim_{\mathbb{R}} \tilde{S}_{n-1}.$$

Whence, we may use Poincaré duality, in order to obtain a surjective map

$$\int : H_c^{b_n+b_{n-1}}(\tilde{S}_{n-1}, \mathbb{C}) \rightarrow \mathbb{C}.$$

To that end, we fix a \mathbb{Q} -valued Haar measure dg_f on $G'(\mathbb{A}_f)$ as follows. For a non-archimedean place $w \in S(\mathcal{K})_f$, we let dg_w be the unique Haar measure on $G'(\mathcal{K}_w)$, which gives $G'(\mathcal{O}_w)$ volume 1 and let dg_f be the product of these local measures. With this choice of a measure, for any open compact subgroup K'_f of $G'(\mathbb{A}_f)$, the volume $\text{vol}(K'_f)$ is a rational number. The set of connected components of the corresponding orbifold $\tilde{S}_{n-1}(K'_f)$ is parameterized by the finite set $\mathcal{X} := \mathbb{A}_f^\times / \mathcal{K}^\times \det(K'_f)$. We assume to have fixed orientations on the various connected components $\tilde{S}_{n-1}(K'_f)_x$, $x \in \mathcal{X}$, given by the orientation on $G'(\mathbb{C})/U(n-1)$, which is defined by the ordered basis $\{X_1, \dots, X_r\}$ of its tangent space at the identity, cf. Sect. 3.3. Hence, the de Rham isomorphism, which is given by integrating over $\tilde{S}_{n-1}(K'_f)_x$ gives rise to a surjective map on $\tilde{S}_{n-1}(K'_f)$,

$$\int_{\tilde{S}_{n-1}(K'_f)} = \sum_{x \in \mathcal{X}} \int_{\tilde{S}_{n-1}(K'_f)_x}.$$

Let $\int_{\tilde{S}_{n-1}}$ stand for the direct limit (over all open compact subgroups K'_f of $G'(\mathbb{A}_f)$) of the normalized maps $\text{vol}(K'_f) \cdot \int_{\tilde{S}_{n-1}(K'_f)}$ and observe that the trivial character $\mathbf{1}$ of $G'(\mathbb{A}_{\mathcal{K}})$ defines a non-trivial cohomology class in $H^0(\tilde{S}_{n-1}, \mathbb{C})$. Using this class, the Poincaré–Duality pairing of $H_c^{b_n+b_{n-1}}(\tilde{S}_{n-1}, \mathbb{C})$ and $H^0(\tilde{S}_{n-1}, \mathbb{C})$ finally gives rise to a surjection

$$(3.6) \quad \begin{array}{ccc} H_c^{b_n+b_{n-1}}(\tilde{S}_{n-1}, \mathbb{C}) & \longrightarrow & \mathbb{C} \\ \theta & \longmapsto & \int_{\tilde{S}_{n-1}} \theta. \end{array}$$

3.9. A non-archimedean, particular vector. We will now choose a special vector in the product of Whittaker models $W(\Pi_f) \times W(\Pi'_f)$, which has the property that it transforms nicely, when plugged into our diagram. This vector will be fixed as in Raghuram [41], 3.1.4, which is itself inspired by Mahnkopf [36], 2.1.1.

First, we remark that for $w \in S(\mathcal{K})_f$, Π'_w is tempered. Hence, Jacquet–Shalika [30], (3.2)

Proposition still holds for Π'_w . Therefore, any non-zero Whittaker functional $\xi'_w \in W(\Pi'_w)$ is non-vanishing on $T'(\mathcal{K}_w)^+ = \{t \in T'(\mathcal{K}_w) \mid t_i t_{i+1}^{-1} \in \mathcal{O}_w, t_{n-1, n-1} = 1\}$. We let $K(m_w)$ (resp. $K'(m'_w)$) be the mirahoric subgroup of $G(\mathcal{K}_w)$ (resp. $G'(\mathcal{K}_w)$) of level m_w (resp. m'_w). This is the subgroup of $G(\mathcal{O}_w)$ (resp. $G'(\mathcal{O}_w)$), consisting of those matrices, whose last row is congruent to $(0, \dots, 0, \star)$ modulo $\wp_w^{m_w}$, where \wp_w is the unique maximal ideal in \mathcal{O}_w . Suppose from now on that m_w (resp. m'_w) is the conductor of Π_w (resp. Π'_w). Then, by Jacquet–Piatetski-Shapiro–Shalika, [28] (5.1) Theorem, the space of Whittaker vectors, transforming by the central character ω_{Π_w} of Π_w (resp. $\omega_{\Pi'_w}$ of Π'_w) under the action of the mirahoric subgroup is one-dimensional and its elements are usually called new vectors.

Now, for Π'_w fix an element $t_{\Pi'_w} \in T'(\mathcal{K}_w)^+$ on which the nontrivial new vectors of Π'_w do not vanish. Observe that we may choose the same element for all σ -twists, i.e., such that $t_{\Pi'_w} = t_{\sigma\Pi'_w}$. If $w \notin S(\Pi')$, then we may take $t_{\Pi'_w} := id$. Depending on these choices, for all $w \in S(\mathcal{K})_f$, we let

$$\xi_{\Pi'_w} := \text{the unique new vector such that } \xi_{\Pi'_w}(t_{\Pi'_w}) = 1$$

This pins down a special Whittaker vector $\xi_{\Pi'_f} := \otimes'_{w \in S(\mathcal{K})_f} \xi_{\Pi'_w} \in W(\Pi'_f)$.

Our choice for Π_w will depend on the data fixed for Π'_w . First, we fix an element $t_{\Pi_w} \in T(\mathcal{K}_w)^+$ analogously as for $G'(\mathcal{K}_w)$. Now, for $w \notin S := S(\Pi') \cup S(\psi)$, we let ξ_{Π_w} be the unique new vector of Π_w , which satisfies $\xi_{\Pi_w}(t_{\Pi_w}) = 1$. It is a certain non-zero multiple c_{Π_w} of the essential vector, cf. [28] (4.1) Théorème. If $w \in S_f$, we take ξ_{Π_w} to be the unique Whittaker vector, whose restriction to $\iota(G'(\mathcal{K}_w))$ is supported on $N'(\mathcal{K}_w)t_{\Pi'_w}K'(m'_w)$ and there equal to $\psi_w^{-1}\omega_{\Pi'_w}^{-1}$. This gives a special Whittaker vector $\xi_{\Pi_f} := \otimes'_{w \in S(\mathcal{K})_f} \xi_{\Pi_w} \in W(\Pi_f)$.

Lemma 3.7. *Let ξ_{Π_f} and $\xi_{\Pi'_f}$ be the above Whittaker vectors. For a non-archimedean place w of \mathcal{K} , the integral*

$$\Psi(s, \xi_{\Pi_w}, \xi_{\Pi'_w}) = \int_{N'(\mathcal{K}_w) \backslash G'(\mathcal{K}_w)} \xi_{\Pi_w}(\iota(g)) \xi_{\Pi'_w}(g) \|\det(g)\|_w^{s-\frac{1}{2}} dg_w$$

converges for $\text{Re}(s) \geq 1 - \mathfrak{m}$ and has a meromorphic continuation to all of \mathbb{C} . It equals

$$\Psi(s, \xi_{\Pi_w}, \xi_{\Pi'_w}) = \begin{cases} c_{\Pi_w} \cdot L(s, \Pi_w \times \Pi'_w) & \text{if } w \notin S \\ \|\det(t_{\Pi'_w})\|_w^{s-\frac{1}{2}} \cdot \text{vol}(K'(m'_w)) & \text{if } w \in S, \end{cases}$$

For all $\sigma \in \text{Aut}(\mathbb{C})$, ${}^\sigma(\xi_{\Pi'_w}) = \xi_{\sigma\Pi'_w}$ for all non-archimedean places w of \mathcal{K} , whereas

$${}^\sigma(\xi_{\Pi_w}) = \begin{cases} \xi_{\sigma\Pi_w} & \text{if } w \notin S \\ \omega_{\sigma\Pi'_w}(t_\sigma) \cdot \xi_{\sigma\Pi_w} & \text{if } w \in S. \end{cases}$$

Proof. Any one of the above assertions is either well-known or follows from a direct calculation using the definition of the local Whittaker vectors. \square

3.10. An archimedean non-vanishing result. Suppose from now on that dg_∞ is the unique Haar measure on $G'(\mathbb{C})$, which gives $U(n-1)$ volume 1. Recall our choices of cohomology classes $[\Pi_\infty]$ and $[\Pi'_\infty]$ from Sect. 3.3. Since $\xi_{\Pi_\infty, i, \alpha}$ is K -finite and $\xi_{\Pi'_\infty, j, \beta}$ is K' -finite, the integral

$$\Psi(s, \xi_{\Pi_\infty, \underline{i}, \alpha}, \xi_{\Pi'_\infty, \underline{j}, \beta}) = \int_{N'(\mathbb{C}) \backslash G'(\mathbb{C})} \xi_{\Pi_\infty, \underline{i}, \alpha}(\iota(g)) \xi_{\Pi'_\infty, \underline{j}, \beta}(g) \|\det(g)\|_\infty^{s-\frac{1}{2}} dg_\infty$$

converges for $\operatorname{Re}(s) \gg 0$. By [8], Thm. 1.2.(i) $\Psi(s, \xi_{\Pi_\infty, \underline{i}, \alpha}, \xi_{\Pi'_\infty, \underline{j}, \beta})$ is holomorphic at all critical values of $L(s, \Pi \times \Pi')$. Assume that $\frac{1}{2} \in \operatorname{Crit}(\Pi \times \Pi')$. In this case, we define

$$c(\tfrac{1}{2}, \Pi_\infty, \Pi'_\infty) := \sum_{\alpha, \beta} \sum_{\underline{i}, \underline{j}} s(\underline{i}, \underline{j}) \cdot \mathcal{T}(e_\alpha \otimes e'_\beta) \cdot \Psi(\tfrac{1}{2}, \xi_{\Pi_\infty, \underline{i}, \alpha}, \xi_{\Pi'_\infty, \underline{j}, \beta}),$$

where $s(\underline{i}, \underline{j})$ is given by $\iota(X_{\underline{i}}^*) \wedge p(X_{\underline{j}}'^*) = s(\underline{i}, \underline{j}) \cdot X_1^* \wedge \dots \wedge X_r^*$, $r = \dim_{\mathbb{R}} \tilde{S}_{n-1}$. By what we said above, $c(\frac{1}{2}, \Pi_\infty, \Pi'_\infty)$ is well-defined, i.e., finite.

Now, drop the assumption that $\frac{1}{2} \in \operatorname{Crit}(\Pi \times \Pi')$ and let $s = \frac{1}{2} + m \in \operatorname{Crit}(\Pi \times \Pi')$ be an arbitrary critical point of $L(s, \Pi \times \Pi')$ with $m \geq 0$. Then, $\frac{1}{2} \in \operatorname{Crit}((\Pi \|\cdot\|^m) \times \Pi')$ and we define

$$c(\tfrac{1}{2} + m, \Pi_\infty, \Pi'_\infty) := c(\tfrac{1}{2}, \Pi_\infty \|\cdot\|^m, \Pi'_\infty).$$

By our compatible choice of $\mathcal{T}^{(m)}$, this is well-defined. The following theorem recently proved by Binyong Sun [46, Thm. A] is of the highest importance for the present paper:

Theorem 3.8. *For all $s = \frac{1}{2} + m \in \operatorname{Crit}(\Pi \times \Pi')$ with $m \geq 0$, $c(\frac{1}{2} + m, \Pi_\infty, \Pi'_\infty) \neq 0$.*

Proof. Since Sun's theorem concerns properties of a non-trivial pairing $\operatorname{Hom}_{G'(\mathbb{C})}(\Pi_\infty \otimes \Pi'_\infty, \mathbb{C})$, and makes no reference to zeta integrals, Whittaker functions, or coefficient systems, we explain how to translate his theorem into our non-vanishing statement. We return to the notation of 3.3. Write $\mathfrak{p} = \mathfrak{g}/\mathfrak{k}$ and $\mathfrak{p}' = \mathfrak{g}'/\mathfrak{k}'$. The expressions there for the generators $[\Pi_\infty] \in H^{b_n}(\mathfrak{g}, K, W(\Pi_\infty) \otimes E_\mu)$ and $[\Pi'_\infty] \in H^{b_{n-1}}(\mathfrak{g}', K', W(\Pi'_\infty) \otimes E_\lambda)$ are based on the following identifications:

$$H_n := H^{b_n}(\mathfrak{g}, K, W(\Pi_\infty) \otimes E_\mu) = \operatorname{Hom}_K(\wedge^{b_n} \mathfrak{p} \otimes E_\mu^*, W(\Pi_\infty)),$$

$$H_{n-1} := H^{b_{n-1}}(\mathfrak{g}', K', W(\Pi'_\infty) \otimes E_\lambda) = \operatorname{Hom}_{K'}(\wedge^{b_{n-1}} \mathfrak{p}' \otimes E_\lambda^*, W(\Pi'_\infty)).$$

Letting h_n, h_{n-1} denote generators of the one-dimensional spaces H_n and H_{n-1} , respectively, it is known that $\rho = \operatorname{Im}(h_n)$ (resp. $\rho' = \operatorname{Im}(h_{n-1})$) is an irreducible K -type in $W(\Pi_\infty)$ (resp. K' -type in $W(\Pi'_\infty)$). Sun's theorem states that, if u is a generator of the one-dimensional space $\operatorname{Hom}_{G'(\mathbb{C})}(\overline{W(\Pi_\infty) \otimes W(\Pi'_\infty)}, \mathbb{C})$, where $\overline{W(\Pi_\infty) \otimes W(\Pi'_\infty)}$ denotes the corresponding Casselman-Wallach (Frechet) completion, then $u|_{\rho \otimes \rho'} \neq 0$. This implies in particular that the Rankin-Selberg integral

$$\Psi_m(\xi \otimes \xi') := \Psi(\tfrac{1}{2} + m, \xi, \xi') = \left[\int_{N'(\mathbb{C}) \backslash G'(\mathbb{C})} \xi(\iota(g)) \xi'(g) \|\det(g)\|_\infty^{s-\frac{1}{2}} dg_\infty \right]_{s=\frac{1}{2}+m}$$

defines a non-zero linear form on $\rho \otimes \rho'$. Here we are using the fact, due to Jacquet, Shalika, and Piatetski-Shapiro, that the Rankin-Selberg integral is not identically zero.

Now if we let $\{e_\alpha^*\}$ and $\{e'_\beta{}^*\}$ denote the dual bases to the bases $\{e_\alpha\}$ and $\{e'_\beta\}$ of E_μ and E_λ introduced in 3.3, we find that (in the obvious notation)

$$\xi_{\Pi_\infty, \underline{i}, \alpha} = h_n(X_{\underline{i}} \otimes e_\alpha^*) \in \rho \quad \xi_{\Pi'_\infty, \underline{j}, \beta} = h_{n-1}(X_{\underline{j}} \otimes e'_\beta{}^*) \in \rho'$$

are the matrix coefficients of h_n and h_{n-1} in the chosen bases. Then the non-vanishing of $c(\frac{1}{2} + m, \Pi_\infty, \Pi'_\infty)$ comes down to the non-vanishing of Ψ_m on $\rho \otimes \rho'$. \square

Observe that Theorem 3.8 implies that $c(\frac{1}{2} + m, \sigma\Pi_\infty, \sigma\Pi'_\infty) \neq 0$, for all $\sigma \in \text{Aut}(\mathbb{C})$. We denote by $p(m, \sigma\Pi_\infty, \sigma\Pi'_\infty)$ the inverse of $c(\frac{1}{2} + m, \sigma\Pi_\infty, \sigma\Pi'_\infty)$. In the special case when $m = 0$, we will abbreviate $p(\sigma\Pi_\infty, \sigma\Pi'_\infty) := p(0, \sigma\Pi_\infty, \sigma\Pi'_\infty)$.

3.11. A theorem on Whittaker periods. Let us recall the Gauß-sum of a Hecke character $\omega : \mathbb{Q}^\times \backslash \mathbb{A}_\mathbb{Q}^\times \rightarrow \mathbb{C}^\times$. Let \mathfrak{c} stand for the conductor ideal of ω_f . We take $y = (y_p)_{p \in S_f} \in \mathbb{A}_f^\times$ such that $\text{ord}_p(y_p) = -\text{ord}_p(\mathfrak{c})$. Then, the Gauß sum of ω_f is defined as $\mathcal{G}(\omega_f, \psi_f, y) = \prod_{p \in S_f} \mathcal{G}(\omega_p, \psi_p, y_p)$, where the local Gauß sum $\mathcal{G}(\omega_p, \psi_p, y_p)$ is defined as

$$\mathcal{G}(\omega_p, \psi_p, y_p) = \int_{\mathbb{Z}_p^\times} \omega_p(u_p)^{-1} \psi_p(y_p u_p) du_p.$$

If all inputs are unramified at p , then $\mathcal{G}(\omega_p, \psi_p, y_p) = 1$, so the above product is finite. Suppressing the dependence on ψ and y , we denote $\mathcal{G}(\omega_f, \psi_f, y)$ simply by $\mathcal{G}(\omega_f)$.

If $\omega : \mathcal{K}^\times \backslash \mathbb{A}_\mathcal{K}^\times \rightarrow \mathbb{C}^\times$ is a Hecke character of \mathcal{K} , let ω_0 denote its restriction to the idèles of \mathbb{Q} . We are now ready to state and prove the main result of this section, which is analogous to the main result in [41].

Theorem 3.9. *Let $\Pi = BC(\pi) \|\cdot\|^m$ be a cuspidal automorphic representation of $G(\mathbb{A}_\mathcal{K})$, $m \in \mathbb{Z}$, and let $\Pi' = BC(\pi')$ be an abelian automorphic representation of $G'(\mathbb{A}_\mathcal{K})$ obtained by base change from unitary groups as in Sections 2.4 and 2.5. In particular, Π is cohomological with respect to E_μ and Π' is cohomological with respect to E_λ . We assume that the highest weights μ and λ satisfy hypothesis 2.3. In view of the archimedean non-vanishing result, cf. Theorem 3.8, the following holds:*

(1) *For all critical values $\frac{1}{2} + m \in \text{Crit}(\Pi \times \Pi')$ with $m \geq 0$ and every $\sigma \in \text{Aut}(\mathbb{C})$,*

$$\sigma \left(\frac{L(\frac{1}{2} + m, \Pi_f \times \Pi'_f)}{p(\Pi)p(\Pi')p(m, \Pi_\infty, \Pi'_\infty)\mathcal{G}(\omega_{\Pi'_f,0})} \right) = \frac{L(\frac{1}{2} + m, \sigma\Pi_f \times \sigma\Pi'_f)}{p(\sigma\Pi)p(\sigma\Pi')p(m, \sigma\Pi_\infty, \sigma\Pi'_\infty)\mathcal{G}(\omega_{\sigma\Pi'_f,0})}.$$

(2)

$$L(\frac{1}{2} + m, \Pi_f \times \Pi'_f) \sim_{\mathbb{Q}(\Pi_f)\mathbb{Q}(\Pi'_f)} p(\Pi)p(\Pi')p(m, \Pi_\infty, \Pi'_\infty)\mathcal{G}(\omega_{\Pi'_f,0}),$$

where “ $\sim_{\mathbb{Q}(\Pi_f)\mathbb{Q}(\Pi'_f)}$ ” means up to multiplication by an element in the composition of number fields $\mathbb{Q}(\Pi_f)\mathbb{Q}(\Pi'_f)$.

Remark 3.10. In fact, the central character of Π' is the twisted base change of the central character of π' , and therefore its restriction to the idèles of \mathbb{Q} is trivial! The term $\mathcal{G}(\omega_{\sigma\Pi'_f,0})$ is nevertheless retained in the statement of the above theorem because it applies, with minor modifications, to cohomological automorphic representations not obtained by base change, in which case the presence of the Gauß sum, as in [41], is indispensable.

Proof. Since (1) implies (2) by Strong Multiplicity One for isobaric automorphic representations, we only prove (1). We will proceed in two steps.

Step 1: Assume $\frac{1}{2}$ is critical.

For any i, j, α and β , denote

$$\varphi_{\Pi, i, \alpha} := W^{-1}(\xi_{\Pi_\infty, i, \alpha} \otimes \xi_{\Pi_f}) \in \Pi \quad \text{and} \quad \varphi_{\Pi', j, \beta} := W^{-1}(\xi_{\Pi'_\infty, j, \beta} \otimes \xi_{\Pi'_f}) \in \Pi',$$

i.e., the inverse image of our particular Whittaker vectors. It follows directly from the definition of the various maps in our diagram, that if we chase $\xi_{\Pi_f} \times \xi_{\Pi'_f}$ through the diagram, we obtain

$$\text{Dia}(\xi_{\Pi_f} \times \xi_{\Pi'_f}) = \sum_{\alpha, \beta} \sum_{\underline{i}, \underline{j}} p(\Pi)^{-1} p(\Pi')^{-1} \cdot s(\underline{i}, \underline{j}) \cdot \mathcal{T}(e_\alpha \otimes e'_\beta) \cdot \int_{G'(\mathcal{K}) \backslash G'(\mathbb{A}_{\mathcal{K}})} \varphi_{\Pi, \underline{i}, \alpha} |_{G'(\mathbb{A}_{\mathcal{K}})} \cdot \varphi_{\Pi', \underline{j}, \beta} dg,$$

where $dg = dg_\infty \cdot dg_f$ (cf. Sect. 3.8 and 3.10). We are left to compute the latter integral. Since $\varphi_{\Pi, \underline{i}, \alpha}$ is cuspidal, it is well-known that

$$\int_{G'(\mathcal{K}) \backslash G'(\mathbb{A}_{\mathcal{K}})} \varphi_{\Pi, \underline{i}, \alpha}(\iota(g)) \cdot \varphi_{\Pi', \underline{j}, \beta}(g) \|\det(g)\|^{s-\frac{1}{2}} dg$$

converges for all $s \in \mathbb{C}$ and equals

$$\int_{N'(\mathbb{A}_{\mathcal{K}}) \backslash G'(\mathbb{A}_{\mathcal{K}})} (\xi_{\Pi_\infty, \underline{i}, \alpha} \otimes \xi_{\Pi_f})(\iota(g)) \cdot (\xi_{\Pi'_\infty, \underline{j}, \beta} \otimes \xi_{\Pi'_f})(g) \|\det(g)\|^{s-\frac{1}{2}} dg$$

for $\text{Re}(s) \gg 0$. For such s , this is furthermore equal to

$$\Psi(s, \xi_{\Pi_\infty, \underline{i}, \alpha}, \xi_{\Pi'_\infty, \underline{j}, \beta}) \cdot L^S(s, \Pi \times \Pi') \prod_{w \notin S} c_{\Pi_w} \prod_{w \in S_f} \|\det(t_{\Pi_w})\|_w^{s-\frac{1}{2}} \text{vol}(K'(m'_w))$$

with $S = S(\Pi') \cup S(\psi)$ by Lemma 3.7. By analytic continuation, we obtain

$$\text{Dia}(\xi_{\Pi_f} \times \xi_{\Pi'_f}) = \frac{L(\frac{1}{2}, \Pi_f \times \Pi'_f)}{p(\Pi)p(\Pi')p(\Pi_\infty, \Pi'_\infty)} \cdot \frac{\prod_{w \notin S} c_{\Pi_w} \prod_{w \in S_f} \text{vol}(K'(m'_w))}{\prod_{w \in S} L(\frac{1}{2}, \Pi_w \times \Pi'_w)},$$

so

$$\sigma(\text{Dia}(\xi_{\Pi_f} \times \xi_{\Pi'_f})) = \sigma \left(\frac{L(\frac{1}{2}, \Pi_f \times \Pi'_f)}{p(\Pi)p(\Pi')p(\Pi_\infty, \Pi'_\infty)} \right) \cdot \frac{\prod_{w \notin S} \sigma(c_{\Pi_w}) \prod_{w \in S_f} \text{vol}(K'(m'_w))}{\prod_{w \in S_f} \sigma(L(\frac{1}{2}, \Pi_w \times \Pi'_w))}.$$

Here we note that by our choice of a measure, cf. Sect. 3.8, $\prod_{w \in S_f} \text{vol}(K'(m'_w))$ is a rational number. Next, observe that $\sigma(L(\frac{1}{2}, \Pi_w \times \Pi'_w)) = L(\frac{1}{2}, {}^\sigma \Pi_w \times {}^\sigma \Pi'_w)$, which is proved in Raghuram [41], Prop. 3.17. Moreover, since the results of Jacquet–Piatetski-Shapiro–Shalika in [29] and [28] as well as Clozel [6] Lemme 4.6 are valid for Π_w , $w \notin S(\psi)$, the proof of Mahnkopf [37] Prop. 2.3.(c) may be carried over to the situation considered here. In other words, $\sigma(c_{\Pi_w}) = c_{{}^\sigma \Pi_w}$ for all $w \notin S$, cf. [36] p. 621 or [41] Prop. 3.21. Finally, we obtain

$$\sigma(\text{Dia}(\xi_{\Pi_f} \times \xi_{\Pi'_f})) = \sigma \left(\frac{L(\frac{1}{2}, \Pi_f \times \Pi'_f)}{p(\Pi)p(\Pi')p(\Pi_\infty, \Pi'_\infty)} \right) \cdot \frac{\prod_{w \notin S} c_{{}^\sigma \Pi_w} \prod_{w \in S_f} \text{vol}(K'(m'_w))}{\prod_{w \in S_f} L(\frac{1}{2}, {}^\sigma \Pi_w \times {}^\sigma \Pi'_w)}.$$

On the other hand, as all maps in the definition of our diagram are σ -equivariant, we see $\sigma(\text{Dia}(\xi_{\Pi_f} \times \xi_{\Pi'_f})) = \text{Dia}({}^\sigma \xi_{\Pi_f} \times {}^\sigma \xi_{\Pi'_f})$. Therefore, Lem. 3.7 implies

$$\sigma(\text{Dia}(\xi_{\Pi_f} \times \xi_{\Pi'_f})) = \left(\prod_{w \in S_f} \omega_{{}^\sigma \Pi'_w}(t_\sigma) \right) \cdot \text{Dia}(\xi_{{}^\sigma \Pi_f} \times \xi_{{}^\sigma \Pi'_f}).$$

As $\omega_{{}^\sigma \Pi'_w}$ is unramified outside $S({}^\sigma \Pi')$ and since $t_\sigma \in \widehat{\mathcal{O}}^*$, we get $\omega_{{}^\sigma \Pi'_w}(t_\sigma) = 1$ for $w \notin S({}^\sigma \Pi')$. Hence, observing that $S(\Pi') = S({}^\sigma \Pi')$, the finite product equals $\prod_{w \in S_f} \omega_{{}^\sigma \Pi'_w}(t_\sigma) =$

$\omega_{\sigma\Pi'_f}(t_\sigma)$. We claim that there is the identity

$$\omega_{\sigma\Pi'_f}(t_\sigma) = \frac{\sigma(\mathcal{G}(\omega_{\Pi'_{f,0}}))}{\mathcal{G}(\omega_{\sigma\Pi'_{f,0}})}.$$

Indeed, at a place $p \in S_f$,

$$\begin{aligned} \mathcal{G}(\omega_{\sigma\Pi'_{p,0}}) &= \int_{\mathbb{Z}_p^\times} \omega_{\sigma\Pi'_{p,0}}(u_p)^{-1} \psi_p(y_p u_p) du_p \\ &= \int_{\mathbb{Z}_p^\times} \sigma(\omega_{\Pi'_{p,0}}(u_p))^{-1} \sigma(\psi_p(y_p u_p t_\sigma^{-1})) du_p \\ &= \sigma(\omega_{\Pi'_{p,0}}(t_\sigma))^{-1} \int_{\mathbb{Z}_p^\times} \sigma(\omega_{\Pi'_{p,0}}(u_p)^{-1} \psi_p(y_p u_p)) du_p \\ &= \omega_{\sigma\Pi'_{p,0}}(t_\sigma)^{-1} \cdot \sigma\left(\int_{\mathbb{Z}_p^\times} \omega_{\Pi'_{p,0}}(u_p)^{-1} \psi_p(y_p u_p) du_p\right) \\ &= \omega_{\sigma\Pi'_{p,0}}(t_\sigma)^{-1} \cdot \sigma(\mathcal{G}(\omega_{\Pi'_{p,0}})), \end{aligned}$$

where the second last equation follows from the fact that – by the very choice of ψ – the above integral is the finite sum over the classes modulo the conductor ideal of $\omega_{\Pi'_p}$ (on which the integrand is constant). Therefore,

$$\sigma\left(\frac{L(\frac{1}{2}, \Pi_f \times \Pi'_f)}{p(\Pi)p(\Pi')p(\Pi_\infty, \Pi'_\infty)}\right) \cdot \frac{\prod_{w \notin S} c_{\sigma\Pi_w} \prod_{w \in S_f} \text{vol}(K'(m'_w))}{\prod_{w \in S_f} L(\frac{1}{2}, \sigma\Pi_w \times \sigma\Pi'_w)} = \frac{\sigma(\mathcal{G}(\omega_{\Pi'_{f,0}}))}{\mathcal{G}(\omega_{\sigma\Pi'_{f,0}})} \cdot \text{Dia}(\xi_{\sigma\Pi_f} \times \xi_{\sigma\Pi'_f}).$$

Recalling that $S(\Pi') = S(\sigma\Pi')$ and m'_w is the same for Π'_w and $\sigma\Pi'_w$, we obtain by what we have seen above, that

$$\text{Dia}(\xi_{\sigma\Pi_f} \times \xi_{\sigma\Pi'_f}) = \frac{L(\frac{1}{2}, \sigma\Pi_f \times \sigma\Pi'_f)}{p(\sigma\Pi)p(\sigma\Pi')p(\sigma\Pi_\infty, \sigma\Pi'_\infty)} \cdot \frac{\prod_{w \notin S} c_{\sigma\Pi_w} \prod_{w \in S_f} \text{vol}(K'(m'_w))}{\prod_{w \in S_f} L(\frac{1}{2}, \sigma\Pi_w \times \sigma\Pi'_w)}.$$

Comparing the last two equations shows (1) if $\frac{1}{2}$ is critical.

Step 2: The general case.

We drop the assumption that $\frac{1}{2}$ is critical now. Let $s = \frac{1}{2} + m$ be an arbitrary critical value of $L(s, \Pi \times \Pi')$, $m \geq 0$. Then $\frac{1}{2}$ is critical for $L(s, (\Pi \|\cdot\|^m) \times \Pi')$ and we are in the situation considered above. Hence, by what we have just observed,

$$\begin{aligned} \sigma\left(\frac{L(\frac{1}{2}, (\Pi \|\cdot\|^m)_f \times \Pi'_f)}{p(\Pi \|\cdot\|^m)p(\Pi')p(\Pi_\infty \|\cdot\|^m, \Pi'_\infty) \mathcal{G}(\omega_{\Pi'_{f,0}})}\right) &= \frac{L(\frac{1}{2}, \sigma(\Pi \|\cdot\|^m)_f \times \sigma\Pi'_f)}{p(\sigma(\Pi \|\cdot\|^m))p(\sigma\Pi')p(\sigma(\Pi_\infty \|\cdot\|^m), \sigma\Pi'_\infty) \mathcal{G}(\omega_{\sigma\Pi'_{f,0}})} \\ &= \frac{L(\frac{1}{2} + m, \sigma\Pi_f \times \sigma\Pi'_f)}{p(\sigma\Pi \|\cdot\|^m)p(\sigma\Pi')p(m, \sigma\Pi_\infty, \sigma\Pi'_\infty) \mathcal{G}(\omega_{\sigma\Pi'_{f,0}})}. \end{aligned}$$

By [42], Thm. 4.1, the period for Π_f satisfies the relation

$$\sigma\left(\frac{p(\Pi \|\cdot\|^m)}{p(\Pi)}\right) = \frac{p(\sigma\Pi \|\cdot\|^m)}{p(\sigma\Pi)},$$

where we used that $\mathcal{G}(\|\cdot\|_f^n) = 1$. This proves the theorem. \square

Remark 3.11. At no time in the proof did we use the assumption that Π' is of abelian type, other than to avoid introducing additional notation. It is therefore clear that our result also holds for cuspidal automorphic representations Π' of $G'(\mathbb{A}_{\mathcal{K}})$; in other words, the proof of Thm. 3.9 also yields Thm. 1.3. As mentioned in the introduction, in this case, i.e., if Π' is cuspidal, it has recently also been proved by Raghuram in [40], Thm. 1.1 – even over any number field. Moreover, it is clear that we did not need that Π and Π' are constructed via base change, in order to prove Thm. 3.9. However, for later use, and to simplify the determination of the set of critical points, we already assumed this to be the case.

Remark 3.12. Let us finally also remark, that we did not need to fix some open compact subgroup $K_f \subset G(\mathbb{A}_f)$ within the entire process of showing Thm. 3.9.

4. MOTIVES – AN INTERLUDE

This section reviews and extends the results of [21] and [24] that are needed to prove the main results of this paper. The crucial definitions and calculations are scattered in different parts of the two papers in question, allowing space for several confusing sign changes, and their conventions are moreover not quite compatible. We hope that the present section will provide a more convenient reference. Moreover, the results of the earlier papers were proved under unnecessarily restrictive hypotheses; here we have striven for maximum generality, always assuming that the base field is an imaginary quadratic field.

The main purpose of this section is to compare the results on special values proved by automorphic methods to Deligne’s conjecture, stated in the setting of motives for absolute Hodge cycles. This setting is partially hypothetical. It is not known, for example, that two geometric realizations in the cohomology of Shimura varieties of the Galois representations attached to a (motivic) algebraic Hecke character are isomorphic as motives, and in particular that they define the same periods. We are therefore led to introduce automorphic analogues of the invariants that arise in the (motivic) calculation of Deligne’s periods over imaginary quadratic fields. See also [50] for calculations of Deligne periods of motives over \mathbb{Q} .

From now on we let $m = 0$, i.e., the cuspidal automorphic representation Π is simply the unitary base change from π .

4.1. Tensor products of motives. The L -functions of the automorphic representations Π and Π' of the previous section are conjecturally attached to motives over $M(\Pi)$ and $M(\Pi')$ over \mathcal{K} of ranks n and $n - 1$, respectively, with coefficients in finite, possibly non-trivial extensions

$$\boxed{E(\Pi)/\mathbb{Q}(\Pi_f) \quad \text{and} \quad E(\Pi')/\mathbb{Q}(\Pi'_f).}$$

The passage from Π to $M(\Pi)$ involves a standard shift, thus

$$L(s, M(\Pi)) = L(s + \frac{1-n}{2}, \Pi),$$

whose center of symmetry is at $s = \frac{n}{2}$ rather than at $s = \frac{1}{2}$. One can in any case attach Hodge structures and compatible families of ℓ -adic representations to Π and Π' , pure of weight $n - 1$ and $n - 2$, respectively, so the tensor product $M(\Pi) \otimes_{\mathcal{K}} M(\Pi')$ is pure of weight

$2n - 3$, and the center of symmetry of the functional equation of its L -function is at the point $s = n - 1$.

More generally, we can take Π' to be an automorphic representation of $GL_{n'}(\mathbb{A}_{\mathcal{K}})$. We will be particularly interested in the cases $n' = n - 1$ and $n' = 1$, but the methods of the present paper allow us to say quite a lot about the general case. In particular, the center of symmetry of the functional equation is the point $\frac{n+n'-1}{2}$, which is not necessarily an integer; however, we will also be interested in critical values to the right of the center of symmetry. The fact that Π and Π' arise by base change from unitary groups is reflected in the polarization property

$$(4.1) \quad M(\Pi)^c \xrightarrow{\sim} M(\Pi)^\vee(1-n) \quad \text{and} \quad M(\Pi')^c \xrightarrow{\sim} M(\Pi')^\vee(1-n').$$

A formalism relating the various realizations of motives satisfying (4.1) and defining their period invariants is developed in §1.1 of [24], and the Deligne periods of the associated adjoint and tensor product motives are computed in §1.3-1.4 of [24].

Recall that the formula of Thm. 3.9 is valid provided the initial representations π and π' of the definite unitary groups H and H' are cohomological with respect to finite-dimensional representations satisfying the equivalent conditions of Lem. 2.1. Exactly the same conditions appear in (2.3.1) of [24] where they are used to evaluate the Deligne period of $M(\Pi) \otimes_{\mathcal{K}} M(\Pi')$ – here $n' = n - 1$ – when the Ichino-Ikeda conjecture computes the central value $L(\frac{1}{2}, \Pi_f \times \Pi'_f)$ in terms of periods of integrals of automorphic forms on $H \times H'$. This is the same central value that appears on the left-hand side of the formulas in 3.9. We exploit this identification in subsequent sections. Here we complete [24] by determining the set of all critical values and their corresponding Deligne periods, when the inequalities of Lem. 2.1 are satisfied.

Write $M = M(\Pi)$, $M' = M(\Pi')$. As in [24], the Hodge types for M (resp. M') are denoted

$$(4.2) \quad (p_i, q_i) := (p_i, n - 1 - p_i) \quad \text{and} \quad (p_i^c, q_i^c) := (p_i^c, n - 1 - p_i^c) = (q_{n+1-i}, p_{n+1-i})$$

(resp. $(r_j, n' - 1 - r_j)$ and $(r_j^c, n' - 1 - r_j^c)$). Bearing in mind that the cohomological parameters of Π and Π' are respectively given by the tuples μ and λ as in 2.2 (and not by a_i 's and b_j 's as in [24]), we have

$$(4.3) \quad p_i = n - i + \mu_i, \quad r_j = n' - j + \lambda_j$$

For any integer m , the Hodge types for the Tate twist $M(m)$ are

$$(4.4) \quad (p_i - m, n - 1 - p_i - m) \quad \text{and} \quad (p_i^c - m, n - 1 - p_i^c - m) = (n - 1 - p_{n+1-i} - m, p_{n+1-i} - m),$$

where the equality on the right is a consequence of 4.1. We extend the definition of [24] by letting $T(M(m), M')$ be the set of pairs (t, u) of indices, $1 \leq t \leq n$, $1 \leq u \leq n'$, such that $p_t^c - m + r_u^c \geq \frac{w+1}{2}$, and write $T(M, M') = T(M(0), M')$. Using 4.4, we transform the left hand side of this equation to

$$n - 1 - p_{n+1-t} + n' - 1 - r_{n'+1-u} - m \geq \frac{n + n' - 1 - 2m}{2}$$

$$\Leftrightarrow \frac{n + n' - 3}{2} \geq p_{n+1-t} + r_{n'+1-u}$$

and this is true if and only if

$$(4.5) \quad \frac{n + n' + 1}{2} \geq t + u + \mu_{n+1-t} + \lambda_{n'+1-u}.$$

4.2. **Critical values when $n' = n - 1$.** For the moment we let $n' = n - 1$. The tensor product motive we consider is $R(M \otimes M') = R_{\mathcal{K}/\mathbb{Q}}(M \otimes M')$, of weight $w = 2n - 3$. Now in view of 2.1, we make the following

Remark 4.6. When $n' = n - 1$, $T(M, M')$ consists exactly of the set of pairs $\{(t, u) \mid t + u \leq n\}$, of cardinality $\frac{n(n-1)}{2}$.

To calculate the Gamma factor, we follow Serre's recipe (see [9] Sect. 5.2–5.3) and find

$$\prod_{a,b} \Gamma_{\mathbb{C}}(s - p_a - r_b) \times \prod_{t,u} \Gamma_{\mathbb{C}}(s - p_t^c - r_u^c)$$

where the product is taken over pairs such that $p_a + r_b \leq \frac{w-1}{2} = n - 2$ and $p_t^c + r_u^c \leq n - 2$. For the first set we have

$$2n - 1 - a - b + \mu_a + \lambda_b \leq n - 2 \Leftrightarrow n + 1 + \mu_a + \lambda_b \leq a + b.$$

By the branching rules this holds if and only if $a + b \geq n + 1$. For the second set the polarization gives us

$$\begin{aligned} p_t^c + r_u^c &= 2n - 3 - (p_{n+1-t} + r_{n-u}) \leq n - 2 \\ \Leftrightarrow n - 1 &\leq p_{n+1-t} + r_{n-u} \\ \Leftrightarrow n - 1 &\leq t + u - 2 + \mu_{n+1-t} + \lambda_{n-u} \end{aligned}$$

which is true if and only if $n + 1 \leq t + u$.

There is therefore a pole at the integer m unless $m \geq 1 + \sup(\sup_{a+b \geq n+1} p_a + r_b, \sup_{t+u \geq n+1} p_t^c + r_u^c)$. The inequalities 2.1 imply that it suffices to consider the pairs (a, b) and (t, u) with $a + b = n + 1$, $t + u = n + 1$. The minimum is thus

$$m_{min} = 1 + \sup\left(\sup_{a+b=n+1} n - 2 + \mu_a + \lambda_b, \sup_{t+u=n+1} n - 2 - \mu_{n+1-t} - \lambda_{n-u}\right).$$

This gives the lower bound of the critical set, and the functional equation that exchanges s with $w + 1 - s$ implies the upper bound is

$$\begin{aligned} m_{max} &= w + 1 - m_{min} = 2n - 2 - m_{min} \\ &= \inf\left(n - 1 - \sup_{1 \leq a \leq n} (\mu_a + \lambda_{n+1-a}), n - 1 + \inf_{1 \leq u \leq n} (\mu_u + \lambda_{n-u})\right) \\ &= n - 1 + \inf\left(- \sup_{2 \leq a \leq n} (\mu_a + \lambda_{n+1-a}), \inf_{1 \leq u \leq n-1} (\mu_u + \lambda_{n-u})\right). \end{aligned}$$

In the following lemma, $\mu(m)$ denotes the n -tuple $\mu_1 - m \geq \mu_2 - m \geq \dots \geq \mu_n - m$, the highest weight of the representation $E_{\mu}^{\text{unt}} \otimes \det^{-m}$.

Lemma 4.7. *The set of critical points of the L -function $L(s, R(M \otimes M'))$ is the set of integers $s_0 = n - 1 \pm m$, where m runs through the non-negative integers such that $\text{Hom}_{G'(\mathbb{C})}((E_{\mu(m)}^{\text{unt}} \otimes E_{\mu^{\vee}(m)}^{\text{unt}}) \otimes (E_{\lambda}^{\text{unt}} \otimes E_{\lambda^{\vee}}^{\text{unt}}), \mathbb{C}) \neq 0$. The maximum value of m is also the minimum of the distances between successive entries in the sequence of inequalities*

$$\mu_1 \geq -\lambda_{n-1} \geq \mu_2 \geq -\lambda_{n-2} \geq \dots \geq -\lambda_1 \geq \mu_n.$$

Proof. The branching law implies that the existence of $G'(\mathbb{C})$ -equivariant homomorphisms

$$E_{\mu(m)}^{\text{unt}} \otimes E_{\lambda}^{\text{unt}} \otimes \rightarrow \mathbb{C}, \quad E_{\mu^{\vee}(m)}^{\text{unt}} \otimes E_{\lambda^{\vee}}^{\text{unt}} \rightarrow \mathbb{C}$$

is equivalent to the two series of inequalities

$$\begin{aligned} \mu_1 - m &\geq -\lambda_{n-1} \geq \mu_2 - m \geq -\lambda_{n-2} \geq \dots \geq -\lambda_1 \geq \mu_n - m, \\ -\mu_1 - m &\leq \lambda_{n-1} \leq -\mu_2 - m \leq \lambda_{n-2} \leq \dots \leq \lambda_1 \leq -\mu_n - m, \end{aligned}$$

which is equivalent to the determination of m_{max} above. The final sentence is clear. \square

We now derive an expression for $c^+(R(M \otimes M')(m))$, following §1.4 of [24], when $n' = n-1$. Recall the formula in Lemma 1.4.1 of [24]:

$$\begin{aligned} c^+(R(M \otimes M')^\vee) &= c^-(R(M \otimes M')^\vee) \\ &\sim \prod_{(t,u) \in T(M, M')} Q_{n+1-t}(M)^{-1} Q_{n-u}(M')^{-1} \cdot \delta(M \otimes M')^{-1} \\ (4.8) \quad &\sim \prod_{t+u \leq n} Q_{n+1-t}(M)^{-1} Q_{n-u}(M')^{-1} \cdot \delta(M)^{1-n} \cdot \delta(M')^{-n} \end{aligned}$$

Here $\delta(M)$ and $\delta(M')$ are the full period determinants studied in §1.2 of [24], specifically in Lemma 1.2.7, which we recall here. We let $E(M)$ and $E(M')$ be the respective coefficient fields of M and M' .

Lemma 4.9. *There are elements $d(M) \in E(M)$ and $d(M') \in E(M')$ such that*

$$\delta(M)^{-1} \sim d(M)^{\frac{1}{2}} (2\pi i)^{n(n-1)/2} Q_{\det M}^{1/2} \quad \text{and} \quad \delta(M')^{-1} \sim d(M')^{\frac{1}{2}} (2\pi i)^{(n-1)(n-2)/2} Q_{\det M'}^{1/2},$$

where $Q_{\det M} = \prod_{t=1}^n Q_t(M)$ and $Q_{\det M'} = \prod_{u=1}^{n-1} Q_u(M')$.

We introduce the new invariants

$$(4.10) \quad q(M) := d(M)^{\frac{1}{2}} Q_{\det M}^{1/2} \quad \text{and} \quad Q_{\leq r}(M) := \prod_{t \leq r} Q_t(M^c), \quad 1 \leq r \leq n$$

and define $q(M')$ and $Q_{\leq r}(M')$ analogously. Recall Remark 1.3.6 of [24]:

$$Q_t(M^c) \sim Q_{n+1-t}(M)^{-1}, \quad t = 1, \dots, n.$$

Combining this with (4.8), (4.9), (4.10), we obtain

$$\begin{aligned} c^\pm(R(M \otimes M')^\vee) &\sim (2\pi i)^{\frac{n(n-1)^2}{2}} (2\pi i)^{\frac{n(n-1)(n-2)}{2}} \prod_{t=1}^{n-1} Q_t(M^c)^{n-t} \prod_{u=1}^{n-1} Q_u(M'^c)^{n-u} \cdot q(M)^{n-1} q(M')^n \\ &\sim (2\pi i)^{n(n-1)(2n-3)/2} \prod_{r=1}^{n-1} (Q_{\leq r}(M) \cdot Q_{\leq r}(M')) \cdot q(M)^{n-1} q(M')^n \\ (4.11) \quad &\sim (2\pi i)^{n(n-1)(2n-3)/2} \prod_{r=1}^{n-1} [Q_{\leq r}(M) \cdot q(M)] \prod_{r'=1}^{n-2} [Q_{\leq r'}(M') \cdot q(M')] \end{aligned}$$

where we have used the relation $Q_{\leq n-1}(M') \sim q(M')^{-2}$ (cf. [24] Lem. 1.2.7 and [24] (1.2.5)) to obtain the more symmetric expression in the last line. Finally, the polarization gives us an isomorphism for each integer m :

$$R(M \otimes M')(m) \cong R(M \otimes M')^\vee(m + 3 - 2n).$$

We write $P_{\leq r}(M) := Q_{\leq r}(M) \cdot q(M)$, and likewise for M' . It thus follows from [9], Formula (5.1.8) that

$$(4.12) \quad \begin{aligned} c^+(R(M \otimes M')(m)) &\sim (2\pi i)^{n(n-1)(m+3-2n)} c^\pm(R(M \otimes M')^\vee) \\ &\sim (2\pi i)^{\frac{n(n-1)}{2}(2m-2n+3)} \prod_{r=1}^{n-1} P_{\leq r}(M) \prod_{r'=1}^{n-2} P_{\leq r'}(M') \end{aligned}$$

4.3. Critical values when $n' = 1$. We repeat the above calculation but now assume $n' = 1$ and weaken the polarization hypothesis for $M' = M(\Pi')$. We assume that M' is of weight $-\kappa$ and that there is a Dirichlet character α_0 and a non-degenerate pairing as in (1.6.2) of [21]:

$$(4.13) \quad R(M') \otimes R(M') \rightarrow \mathbb{Q}(\alpha_0 \cdot \varepsilon_{\mathcal{K}})(\kappa).$$

We assume moreover that there are Hecke characters χ and α , as in [21] (especially §2.9 and §3.5), so that $\Pi' = BC(\chi) \cdot \alpha$, with $\chi_\infty(z) = z^{-k}$, $\alpha_\infty(z) = z^\kappa$, so that $\Pi'_\infty(z) = (z/\bar{z})^{-k} \cdot z^\kappa$. The motive $R(M')$ then has Hodge types $(k-\kappa, -k); (-k, k-\kappa)$. (The parameter κ is needed for parity considerations and will in practice either be 0 or 1.) The restriction of Π' to the idèles of \mathbb{Q} equals α_0 multiplied by a power of the norm; thus the notation of 4.13 is consistent with that introduced on p. 92 of [21].

The indices b and u only take the value 1; we have $r_1 = k - \kappa$ and $r_1^c = -k$, and the weight of $R(M \otimes M')$ is $n - 1 - \kappa$. By definition $T(M(m), M')$ is the set of pairs $(t, u) = (t, 1)$ such that $p_t^c - m - k \geq \frac{n-\kappa-2m}{2}$.

Remark 4.14. When $n' = 1$, $T(M(m), M')$ consists exactly of the set of pairs $(t, 1)$ such that

$$n - 2p_t^c \leq -2k + \kappa.$$

Since the p_t^c are decreasing, $T(M(m), M')$ is therefore the set of $t \leq s(M')$ where

$$(4.15) \quad \boxed{s(M') := s(M', M) := \sup\{t \mid n - 2p_t^c \leq -2k + \kappa\}}$$

and it is of cardinality $s(M')$. Let

$$\boxed{r(M') := r(M', M) := n - s(M', M)}.$$

If we assume $\kappa \equiv n \pmod{2}$, then, given the relations $p_i^c = n - 1 - p_{n+1-i} = q_{n+1-i}$, we find

$$(4.16) \quad 2k - \kappa \in [n - 2p_{r(M')}, n - 2p_{r(M')+1} - 2] = [n - 2q_{s(M')+1}^c, 2p_{s(M')}^c - n].$$

The set of critical values is determined in [21], (3.3.8.1). When $n > 1$ we choose κ such that the weight w of $R(M) \otimes R(M')$ is odd; then $\frac{w+1}{2}$ is always critical. When $n = n' = 1$ we take $\kappa = 0$ (resp. $\kappa = 1$) and $k \neq 0$; then the near-central point $s = 1$ (resp. the central point $s = 0$) is always critical. As in the previous section, we obtain the formula

$$(4.17) \quad \begin{aligned} c^\pm(R(M \otimes M')^\vee) &\sim \prod_{t \leq s(M')} Q_{n+1-t}(M)^{-1} Q_1(M')^{-1} \cdot \delta(M \otimes M')^{-1} \\ &\sim \prod_{t \leq s(M')} Q_t(M^c) Q_1(M')^{-1} \cdot \delta(M)^{-1} \cdot \delta(M')^{-n} \\ &\sim (2\pi i)^{\frac{n(n-1)}{2}} \cdot P_{\leq s(M')}(M) \cdot Q_1(M')^{-s(M')} \delta(M')^{-n} \end{aligned}$$

The polarizations define an isomorphism

$$R(M \otimes M') \cong R(M \otimes M')^\vee(1 - n + \kappa) \otimes \mathbb{Q}(\alpha_0 \cdot \varepsilon_{\mathcal{K}})$$

and so, as before

$$(4.18) \quad \begin{aligned} & c^+(R(M \otimes M')(m)) \\ & \sim (2\pi i)^{n(1-n+\kappa) + \frac{n(n-1)}{2} + nm} \mathcal{G}(\alpha_{0,f}^{-1} \cdot \varepsilon_{\mathcal{K},f})^n \cdot P_{\leq s(M')}(M) \cdot Q_1(M')^{-s(M')} \delta(M')^{-n} \\ & \sim (2\pi i)^{mn - \frac{n(n-1)}{2}} \mathcal{G}(\alpha_{0,f}^{-1} \cdot \varepsilon_{\mathcal{K},f})^n \cdot P_{\leq s(M')}(M) \cdot (2\pi i)^{n\kappa} Q_1(M')^{-s(M')} \delta(M')^{-n} \end{aligned}$$

4.4. Holomorphic automorphic forms on unitary groups and automorphic critical intervals when $n' = 1$. In [21] and its successor [22], expressions for the critical values of L -functions of unitary groups are derived from the determination of fields of rationality of Eisenstein series for the Siegel parabolic, on the one hand, and from the construction of holomorphic differential operators. The latter are used to specify the signature of the unitary similitude group whose associated Shimura variety realizes the critical values as the Petersson square norm of a (rational) holomorphic automorphic form, up to elementary factors. Unfortunately the definitions of the parameters are scattered in several places with confusing sign changes between successive appearances. For the reader's convenience, the definitions of all relevant parameters and the main results on the existence of differential operators are reproduced here.

Page numbers refer to [21]. The parameter (μ_1, \dots, μ_n) of Π is denoted (a_1, \dots, a_n) , starting on p. 104; we replace the a 's with μ 's in what follows. There is also a central parameter \mathbf{c} (denoted c in [21]; we use boldface to distinguish the parameter from complex conjugation) and assumed (on p. 103) to be of the same parity as $\sum_{i=1}^n \mu_i$, and such that the motives realized on the degree i cohomology of unitary group Shimura varieties are of weight $i - \mathbf{c}$ (see p. 104). The main results of [21] apply to self-dual Π and with our conventions there it follows (though this is not adequately explained) that $\mathbf{c} = 0$. Here we assume for simplicity that $\mathbf{c} = \sum_{i=1}^n \mu_i = C(\mu)$ in the notation of p. 105. This implies that the integers $\mathcal{P}(\mu)$ and $\mathcal{Q}(\mu)$ defined on p. 105 are given by

$$\mathcal{P}(\mu) = -\mathbf{c} \quad \text{and} \quad \mathcal{Q}(\mu) = 0.$$

We fix a signature (r, s) , with $n = r + s$. For any integer k one defines

$$\mu(k) := (\mu_1 - k, \mu_2 - k, \dots, \mu_n - k; -nk + \mathbf{c}),$$

$$\Lambda(\mu(k); r, s) := (\mu_{s+1} - k - s, \dots, \mu_n - k - s; \mu_1 - k + r, \dots, \mu_s - k + r; -nk + \mathbf{c}).$$

(buried on p. 136 between (2.9.8) and (2.9.9), with r and s switched).

The motive $M(\Pi)$ attached abstractly to Π has the Hodge types given by 4.2. These cannot always be realized as cohomological motives, for several reasons. If Π descends to an (L -packet of) automorphic representations π of a unitary group $U(V)$, V is a Hermitian space over \mathcal{K} of signature (r, s) , then we obtain a motive with coefficients in \mathcal{K} , denoted $M'(\pi, V; r)$ on p. 118, in the cohomology of the corresponding Shimura variety, denoted $Sh(V)$ in [21]. The L -packet contains the holomorphic type representation denoted $\pi_{r,s}$ above. When Π is self-dual, as in [21], this is sufficient ², provided there is a unitary group

²More precisely, to get (the Betti realization of) a legitimate motive $M'(\pi, V; r)$ we need to take the sum $M'(\pi, V; r) \oplus M'(\pi, -V; s)$, as on p. 118, and this is necessary to define the archimedean Frobenius as an operator on automorphic forms; but for the de Rham structure it suffices to work with $M'(\pi, V; r)$.

$U(V)$ of signature $(n-1, 1)$ to which Π descends. When n is even there is a local obstruction and one may only be able to realize *even* exterior powers of the desired $M(\Pi)$, even in the self-dual case. We therefore make the following hypothesis:

Hypothesis 4.19. For every signature (r, s) with $r + s = n$, there is a unitary group $U(V)$ of signature (r, s) such that the cuspidal automorphic representation Π of $G(\mathbb{A}_{\mathcal{K}})$ descends to an L -packet of automorphic representations $\{\pi\}$ of $U(V)$. Moreover, the L -packet $\{\pi\}$ contains a member whose archimedean component is in the holomorphic discrete series.

This hypothesis will be assumed implicitly in the remainder of this section. In the statements of the main theorems in Section 6 the hypothesis will be included explicitly; there will also be a discussion of substitutes.

When Π is conjugate self-dual but not self-dual – more precisely, when the central character ω_{Π} of Π is not trivial, in particular when the integer $C(\mu)$ defined above is not even – there may be an additional obstruction. We need to choose a Hecke character ξ such that

$$(4.20) \quad \tilde{\xi} := \xi/\xi^c = \omega_{\Pi}^{-1}.$$

Such a choice is always possible, cf. Proposition 1.2.4 of [7] (which cites the book of Harris and Taylor). This is because the Shimura variety is defined in terms of $GU(V)$ and not $U(V)$, and an automorphic representation π of $GU(V)$ gives rise by base change to a pair of automorphic representations (Π, ξ) of $GL_n(\mathbb{A}_{\mathcal{K}}) \times GL_1(\mathbb{A}_{\mathcal{K}})$. The parameter \mathbf{c} is determined by the archimedean component ξ_{∞} of ξ :

$$\xi_{\infty}(t) = t^{-\mathbf{c}}.$$

The Hodge type of the motive realized in the cohomology of $Sh(V_1)$, with $(r, s) = (n-1, 1)$ (see [21], Example 2.2.5) – call it $M'(\pi, V_1; n-1)$ (and don't confuse this M' with the M' for GL_{n-1}) – is called the *principal Hodge type* in §2.2.9 of [21]; it is given by

$$(4.21) \quad (p_i(\mu), q_i(\mu)) = (p_i + \mathcal{P}(\mu), q_i + \mathcal{Q}(\mu)) = (p_i - \mathbf{c}, q_i),$$

where (p_i, q_i) are attached to Π as in 4.2. Observe that by (4.3), the pairs (p_i, q_i) also depend on μ (although this has been hidden in the notation). More generally, if V_s has signature (r, s) , the motive $M'(\pi_{r,s}, V_s; r)$ attached to the (holomorphic) representation $\pi_{r,s}$ of $GU(V)$ that is nearly equivalent to π , in the sense that π and $\pi_{r,s}$ have the same base change (Π, ξ) to $GL_n(\mathbb{A}_{\mathcal{K}}) \times GL_1(\mathbb{A}_{\mathcal{K}})$, is expected to bear the following relation to the desired $M(\Pi)$:

$$(4.22) \quad M'(\pi_{r,s}, V_s; r) \cong \wedge^r(M(\Pi)^{\vee}) \otimes M(\xi^c)$$

This is known to be true in most cases at the level of Galois representations (cf. [7], Proposition 4.3.8, as well as the more complete results of Shin and Scholze–Shin), but in the absence of the Tate conjecture one can't even give a precise definition to the rank one motive $M(\xi^c)$, much less prove a result such as (4.22). The best we can do is to show how the expressions for special values obtained by analytic methods are products of expressions that resemble those that would be expected if there were a complete theory of motives. The choice of ξ should have no bearing on the expression of the special value of the L -function of Π ; we return to this point below.³

³Actually, the Tate conjecture is more than we need. It would be enough to show that an isomorphism at the level of Galois representations implies an isomorphism at the level of motives for absolute Hodge cycles.

We can now restate Corollary 3.3.8 of [21], which is the main result on the existence of holomorphic differential operators. In what follows, notation is as on p. 145. In particular, $\mathcal{E}_{m,\kappa}$ is a line bundle on the Shimura variety $Sh(n, n)$ (attached to a quasi-split unitary similitude group of size $2n$ and $E_{\Lambda, \Lambda^\sharp(\kappa)}$ is an automorphic vector bundle on the Shimura variety $Sh(V, -V)$ (essentially $Sh(V) \times Sh(-V)$).

Lemma 4.23. *Let $n' = 1$ and Π' , $M' = M(\Pi')$, k , and κ be as in 4.3. Let $(r, s) = (r(M'), s(M'))$ and let V be a Hermitian space of signature (r, s) . Let m be a critical value of $R(M \otimes M')$ to the right of (the central point) $\frac{1}{2}(n - \kappa)$. There is a (non-trivial) differential operator*

$$\Delta(m, \kappa, \Lambda) : \mathcal{E}_{m,\kappa} \rightarrow E_{\Lambda, \Lambda^\sharp(\kappa)}$$

where $\Lambda = \Lambda(\mu^\vee(k); r, s)$ on the Shimura variety $Sh(V, -V)$.

Proof. Keeping in mind that in absence of the self-duality hypothesis for μ we have to replace μ by μ^\vee in corollary 3.3.8 of [21], we obtain such a differential operator for the pair (r, s) , provided

$$\frac{1}{2}(n - \kappa) \leq m \leq \min(q_{s+1}(\mu^\vee) + k - \kappa - \mathcal{Q}(\mu^\vee), p_s(\mu^\vee) - k - \mathcal{P}(\mu^\vee)) = \min(q_{s+1}^c + k - \kappa, p_s^c - k),$$

where the second equality is (4.21) for μ^\vee . In particular, we have $2q_{s+1}^c + 2k \geq n + \kappa$ and $2p_s^c - 2k \geq n - \kappa$; in other words

$$(4.24) \quad 2k - \kappa \in [n - 2q_{s+1}^c, 2p_s^c - n].$$

So (4.16) implies that (4.24) is true exactly when $s = s(M') = s(M', M)$. \square

We now summarize informally the constructions of §3 of [21] and the corresponding sections of [22]. We let α and χ be Hecke characters of $\mathbb{A}_{\mathcal{K}}$ and $U(1)(\mathbb{A})$, respectively, with parameters κ and k as above. Let π be an anti-holomorphic representation of $GU(V)(\mathbb{A})$ and identify its contragredient π^\vee with an anti-holomorphic representation of $GU(-V)(\mathbb{A})$ as in [21] (this identification is spread out over several sections; the most important statements are Corollary 2.5.11.2, Lemma 2.8.8, and (2.8.9.5)). If $Eis \in \Gamma(Sh(n, n), \mathcal{E}_{m,\kappa})$ – in practice a holomorphic Eisenstein series – and $f \in \pi$, $f' \in \pi^\vee$, and $f_\chi = f \otimes \chi \circ \det$, $f'_\chi = f' \otimes \chi^{-1} \circ \det$, then the value of the standard L -function at $s = m$ can be identified, up to elementary and local terms, with the cup product (Serre duality) pairing

$$(4.25) \quad \Delta(m, \kappa, \Lambda) \cup [f_\chi \otimes (f'_\chi \otimes \alpha^{-1} \circ \det)] \rightarrow \mathbb{C}$$

This pairing is well defined provided (in the notation of p. 147)

$$(4.26) \quad f_\chi \otimes (f'_\chi \otimes \alpha^{-1} \circ \det) \in \overline{H}^{2rs}(Sh(V, -V), \Omega_{Sh}^{top} \otimes E_{\Lambda, \Lambda^\sharp(\kappa)}^\vee).$$

where Ω_{Sh}^{top} is the canonical bundle on $Sh(V, -V)$ and \overline{H} is the version of coherent cohomology used in [21]. Now there is an isomorphism of automorphic vector bundles:

$$(4.27) \quad \Omega_{Sh}^{top} \otimes E_{\Lambda, \Lambda^\sharp(\kappa)}^\vee \cong E_{\mu(-k), \mu(k-\kappa)},$$

with $\Lambda = \Lambda(\mu^\vee(k); r, s)$ as in Lem. 4.23. This isomorphism is stated in the third displayed formula on p. 150 of [21] with μ in place of μ^\vee inside Λ , which is legitimate in that setting because of the self-duality hypothesis. Now (4.27) implies that (4.26) holds, since Π_∞ has cohomology with respect to E_μ .

Therefore, the above calculations apply and give us the following version of Theorem 3.5.13 of [21] and Theorem 4.3 of [22]. Recall that when β is a Hecke character, we have

defined $\check{\beta} = (\beta^c)^{-1}$ on p. 82 of [21]. This should not be confused with $\beta^\vee = \beta^{-1}$. In what follows, we let $\pi = \pi(\Pi, \xi)$ be an anti-holomorphic automorphic representation of $GU(V)(\mathbb{A})$, for some Hermitian space V to be specified, whose base change equals (Π, ξ) . For any integer $j \in [0, n]$ define $P^{(j)}(\Pi, \xi)$ to be the period denoted $P^{(j)}(\pi, \star, \beta)$ in [21], (2.8.2): It is the Petersson inner product with itself of a rationally normalized holomorphic form on $GU(V)$ whose restriction to $U(V)$ is weakly equivalent to π on a Hermitian space of signature $(n - j, j)$. We set

$$(4.28) \quad \boxed{P^{(j)}(\Pi) := (2\pi)^c P^{(j)}(\Pi^\vee, \xi^{-1}).}$$

This can be shown to be independent of the choice of ξ when appropriate L -functions have non-vanishing critical values (see 4.33 for an explanation). The twist by the contragredient is used for notational purposes only. We write

$$\boxed{G(\Pi') := (2\pi i)^k p(\check{\chi}^{(2)} \cdot \check{\alpha}, 1).}$$

The notation $\alpha_{f,0}$ for the restriction of (the finite part of) a Hecke character to the idèles of \mathbb{Q} , and \mathcal{G} for the Gauß sum, are as in §3.

Theorem 4.29. *Let $m \geq \frac{1}{2}(n - \kappa)$ be a critical value of $L(s, R(M(\Pi) \otimes M(\Pi')))$. Let $(r, s) = (r(M'), s(M'))$ and let V be a Hermitian space of dimension n over \mathcal{K} with signature (r, s) . Let ξ be a Hecke character of \mathcal{K} satisfying 4.20. Then,*

$$L(m, R(M(\Pi) \otimes M(\Pi'))) \sim_{\mathcal{K}E(\Pi)E(\Pi')} (2\pi i)^{mn - \frac{n(n-1)}{2}} \mathcal{G}(\varepsilon_{\mathcal{K},f})^{\lceil \frac{n}{2} \rceil} P^{(s)}(\Pi^\vee) [(2\pi i)^\kappa \mathcal{G}(\alpha_{f,0}^{-1})]^s G(\Pi')^{n-2s}$$

As mentioned above, this is a restatement of Theorem 3.5.13 of [21] and Theorem 4.3 of [22]. At the suggestion of the referee, we remind the reader that this result is proved by using the doubling method of Piatetski–Shapiro–Rallis and Garrett to identify the L -function with an integral of a degenerate Eisenstein series on a quasi-split unitary group to which the Shimura variety $Sh(n, n)$ is attached, against antiholomorphic forms on the Shimura subvariety $Sh(V, -V)$. The statement concerns critical values to the right of the center of symmetry; these correspond precisely to values of the Eisenstein series that are nearly holomorphic in Shimura’s sense, and also to the differential operators described in 4.23. The critical values can thus be analyzed in terms of cup products in coherent cohomology of $Sh(V, -V)$. Note that the motivic interpretation of these critical values depends on an interpretation involving coherent cohomology, whereas the critical values of the Rankin–Selberg L -function of $GL(n) \times GL(n - 1)$ are interpreted as cup products in topological cohomology.

Thm. 4.29 has the following equivalent reformulation. To this end, we recall that

$$R(M(\Pi) \otimes M(\Pi')) \equiv R(M(\Pi^c) \otimes M(\Pi'^c)) = R(M(\Pi^\vee) \otimes M(\Pi'^c)).$$

Hence, by replacing μ by μ^\vee in (4.27), we may apply Thm. 4.29 to the pair $(M(\Pi^\vee), M(\Pi'^c))$, but still obtain an expression for our original L -value $L(m, R(M(\Pi) \otimes M(\Pi')))$. Let us write

$$\Pi'^c = BC(\chi)^c \alpha^c = BC(\chi') \cdot \alpha', \quad M'^c = M(\Pi'^c).$$

Then, a simple calculation shows that $k' = -k + \kappa$ and $\kappa' = \kappa$, where k' is attached to χ' (resp. κ' to α') as k has been attached to χ (resp. κ to α). Hence, replacing M' by M'^c

in (4.16), we obtain $2k' - \kappa' \in [n - 2q_{s+1}^c(\Pi), 2p_s^c(\Pi) - n]$ if and only if $s = s(M'^c, M)$. Inserting the above values for k' and κ' and inverting the interval, we end up with

$$2k - \kappa \in [n - 2p_s^c(\Pi), 2q_{s+1}^c(\Pi) - n] = [n - 2q_{r+1}(\Pi), 2p_r(\Pi) - n],$$

if and only if $r = r(M'^c, M)$. Now, recall that

$$2k - \kappa \in [n - 2q_{s+1}(\Pi), 2p_s(\Pi) - n] = [n - 2q_{s+1}^c(\Pi^\vee), 2p_s^c(\Pi^\vee) - n]$$

if and only if $s = s(M', M^\vee)$. Thus,

$$(4.30) \quad (r(M'^c, M^\vee), s(M'^c, M^\vee)) = (s(M', M), r(M', M)).$$

We may hence reformulate Thm. 4.29 by

Theorem 4.31. *Let $m \geq \frac{1}{2}(n - \kappa)$ be a critical value of $L(s, R(M(\Pi) \otimes M(\Pi^\vee)))$. Let $(r, s) = (s(M'), r(M'))$ and let V be a Hermitian space of dimension n over \mathcal{K} with signature (r, s) . Let ξ be a Hecke character of \mathcal{K} satisfying 4.20. Then,*

$$L(m, R(M(\Pi) \otimes M(\Pi^\vee))) \sim_{\mathcal{K}E(\Pi)E(\Pi^\vee)} (2\pi i)^{mn - \frac{n(n-1)}{2}} \mathcal{G}(\varepsilon_{\mathcal{K},f})^{\lceil \frac{n}{2} \rceil} P^{(r(M'))}(\Pi) [(2\pi i)^\kappa \mathcal{G}(\alpha_{f,0}^{-1})]^{r(M')} G(\Pi'^c)^{n-2r(M')}$$

4.5. A comparison to Deligne's periods. Now let us compare this to the critical value predicted by Deligne's conjecture, namely (4.18):

$$(2\pi i)^{mn - \frac{n(n-1)}{2}} \mathcal{G}(\alpha_{0,f}^{-1} \cdot \varepsilon_{\mathcal{K},f})^n \cdot P_{\leq s(M')}(M) \cdot (2\pi i)^{n\kappa} Q_1(M')^{-s(M')} \delta(M')^{-n}.$$

Or rather let us make the following

Optimistic Comparison 4.32. Compare

$$(Aut) \quad \mathcal{G}(\varepsilon_{\mathcal{K},f})^{\lceil \frac{n}{2} \rceil} P^{(r(M'))}(\Pi) [(2\pi i)^\kappa \mathcal{G}(\alpha_{0,f}^{-1})]^{r(M')} G(\Pi'^c)^{n-2r(M')}$$

to

$$(Mot) \quad \mathcal{G}(\alpha_{0,f}^{-1} \cdot \varepsilon_{\mathcal{K},f})^n \cdot P_{\leq s(M')}(M) \cdot (2\pi i)^{n\kappa} Q_1(M')^{-s(M')} \delta(M')^{-n}.$$

Applying (4.22), and bearing in mind that the base change of π is (Π, ξ) , we see that $i^c P^{(r(M'))}(\Pi)$ is the correct normalization for the Petersson inner product of a rational class in the bottom stage of the Hodge filtration of $\wedge^{s(M')} M(\Pi^\vee) \otimes M((\xi^c)^{-1})$. (The presence of the power i^c , which will disappear in our applications in any case, is explained in [21], Lemma 2.8.8.) On the other hand, $P_{\leq s(M')}(M) = Q_{\leq s(M')}(M) \cdot q(M)$ is the product with $q(M)$ of the Petersson norm of a rational class in the bottom stage of the Hodge filtration of $\wedge^{s(M')} M(\Pi^\vee)$, and plausible arguments can be made to justify identifying $(2\pi i)^c q(M)$ with the Petersson norm of $M((\xi^c)^{-1})$ (cf. [21], (3.7.9.3)-(3.7.9.5), where however it is assumed that $\mathbf{c} = 0$).

Remark 4.33. Note that the automorphic side (Aut) of the comparison involves a Petersson norm on the unitary similitude group, and therefore depends in principle on the auxiliary Hecke character ξ defined in (4.20). The motivic side (Mot), on the other hand, is independent of this choice. Now Theorem 4.29 shows that replacing ξ with ξ' , also assumed to satisfy (4.20), gives another expression for the same critical value. In particular, if the critical value does not vanish, then we see that $P^{(r(M'))}(\Pi^\vee, \xi) \sim P^{(r(M'))}(\Pi^\vee, \xi')$, where \sim is taken over a simultaneous field of definition for Π, ξ , and ξ' . The "plausible arguments" mentioned above give a motivic interpretation for this identity.

It thus remains to justify an identification

$$(4.34) \quad \mathcal{G}(\alpha_{0,f}^{-1} \cdot \varepsilon_{\mathcal{K},f})^n \cdot (2\pi i)^{n\kappa} Q_1(M')^{-s(M')} \delta(M')^{-n} \sim \mathcal{G}(\varepsilon_{\mathcal{K},f})^{\lceil \frac{n}{2} \rceil} (2\pi i)^{\kappa r(M')} \mathcal{G}(\alpha_{0,f}^{-1})^{r(M')} G(\Pi'^c)^{n-2r(M')}.$$

We provide such a justification, at least up to Galois conjugation over \mathcal{K} , in the next section.

4.6. The case of a Hecke character. We now assume $n = 1$ and $\Pi = \mathbf{1}$ is the trivial character, so $M = \mathbb{Q}(0)$. We have $s(M') = s(M', \mathbb{Q}(0)) = 1$ if $-2k + \kappa \geq 1$, $s(M') = 0$ if $-2k + \kappa \leq 0$. Then, if $R(M')(m)$ is critical, (4.18) becomes

$$(4.35) \quad c^+(R(M')(m)) \sim \begin{cases} (2\pi i)^{m+\kappa} \mathcal{G}(\alpha_{0,f}^{-1} \cdot \varepsilon_{\mathcal{K},f}) \cdot Q_1(M')^{-1} \delta(M')^{-1}, & \text{if } 2k - \kappa \leq -1 \\ (2\pi i)^{m+\kappa} \mathcal{G}(\alpha_{0,f}^{-1} \cdot \varepsilon_{\mathcal{K},f}) \cdot \delta(M')^{-1} & \text{if } 2k - \kappa \geq 0 \end{cases}$$

Since Deligne's conjecture is known for critical values of $L(s, R(M'))$, we thus have

$$(4.36) \quad L(m, R(M')) \sim \begin{cases} (2\pi i)^{m+\kappa} \mathcal{G}(\alpha_{0,f}^{-1} \cdot \varepsilon_{\mathcal{K},f}) \cdot Q_1(M')^{-1} \delta(M')^{-1}, & \text{if } 2k - \kappa \leq -1 \\ (2\pi i)^{m+\kappa} \mathcal{G}(\alpha_{0,f}^{-1} \cdot \varepsilon_{\mathcal{K},f}) \cdot \delta(M')^{-1} & \text{if } 2k - \kappa \geq 0 \end{cases}$$

Here \sim means $\sim_{\mathbb{Q}(M');\mathcal{K}}$ as in [21], because we are going to compare this with the result of Thm. 4.31:

$$(4.37) \quad L(m, R(M')) \sim (2\pi i)^{m+r(M')\kappa} \mathcal{G}(\varepsilon_{\mathcal{K},f}) \mathcal{G}(\alpha_{0,f}^{-1})^{r(M')} G(\Pi'^c)^{1-2r(M')}$$

Assume $s(M') = 0$, so $2k - \kappa \geq 0$. Then comparing (4.37) and (4.36) we find

$$(4.38) \quad \mathcal{G}(\alpha_{0,f}^{-1} \cdot \varepsilon_{\mathcal{K},f}) \cdot \delta(M')^{-1} \sim \mathcal{G}(\varepsilon_{\mathcal{K}}) \mathcal{G}(\alpha_{0,f}^{-1}) G(\Pi'^c)^{-1}.$$

Let us now get back to (4.34). Suppose for the moment $\alpha = 1$, i.e., $\kappa = 0$. So we compare $\mathcal{G}(\varepsilon_{\mathcal{K},f})^n Q_1(M')^{-s(M')} \delta(M')^{-n}$ to $\mathcal{G}(\varepsilon_{\mathcal{K},f})^{\lceil \frac{n}{2} \rceil} G(\Pi'^c)^{n-2r(M')}$. Inserting (4.38), we finally have to compare

$$\mathcal{G}(\varepsilon_{\mathcal{K},f})^n Q_1(M')^{-s(M')} \delta(M')^{-n}$$

to

$$\mathcal{G}(\varepsilon_{\mathcal{K},f})^{\lceil \frac{n}{2} \rceil} \delta(M')^{n-2r(M')}.$$

Now observe that [24] Lem. 1.2.7 and [24] (1.2.5) imply that $Q_1(M') \sim \delta(M')^{-2}$. So we find that

$$\mathcal{G}(\varepsilon_{\mathcal{K},f})^n Q_1(M')^{-s(M')} \delta(M')^{-n} \sim \mathcal{G}(\varepsilon_{\mathcal{K},f})^n \delta(M')^{2s(M')-n} \sim \mathcal{G}(\varepsilon_{\mathcal{K},f})^n \delta(M')^{n-2r(M')}.$$

Hence, the two side of (4.34) coincide up to a power of the Gauß sum $\mathcal{G}(\varepsilon_{\mathcal{K},f})$. We remark that the same conclusion holds if one uses the case $s(M') = 1$ in order to obtain a relation for $G(\Pi'^c)$. Moreover, the case of non-trivial α is only slightly more complicated, and will not be used in the sequel.

5. RATIONAL EISENSTEIN CLASSES OF ABELIAN TYPE

5.1. Boundary cohomology. Recall the abelian representation Π' from Sect. 2.5 and the space S_{n-1} from Sect. 3.1. We denote by \overline{S}_{n-1} the Borel-Serre compactification of S_{n-1} and by $\partial \overline{S}_{n-1}$ its boundary, cf. [3], [43]. The face coming from the Borel subgroup B' of G' is denoted $\partial_{B'} \overline{S}_{n-1}$. It is given by

$$\partial_{B'} \overline{S}_{n-1} = B'(\mathcal{K}) \backslash G'(\mathbb{A}_{\mathcal{K}}) / K'.$$

Since the only automorphic representations of the torus $T' \subset B'$ are its Hecke characters χ it follows that

$$H^q(\partial_{B'} \bar{S}_{n-1}, \mathcal{E}_\lambda) = \bigoplus_{\chi} H^q(\mathfrak{g}', K', \text{Ind}_{B'(\mathbb{A}_{\mathcal{K}})}^{G'(\mathbb{A}_{\mathcal{K}})}[\chi] \otimes E_\lambda),$$

see e.g. Harder [18]. Observe that induction is unitary here and that $\pi_0(G'(\mathbb{C})) = \pi_0(B'(\mathbb{C})) = 0$, which simplifies things. Recall the character $\tilde{\tau} = BC(\chi_1)\gamma \otimes \dots \otimes BC(\chi_{n-1})\gamma$ from Sect. 3.4 and let $W(G')$ be the Weyl group of G' . For $w \in W(G')$, we let $\tilde{\tau}^w$ be the representation

$$\tilde{\tau}^w := BC(\chi_{w^{-1}(1)})\gamma \otimes \dots \otimes BC(\chi_{w^{-1}(n-1)})\gamma$$

of $T'(\mathbb{A}_{\mathcal{K}})$, where w is viewed as a permutation. In particular, we may write

$$H^q(\partial_{B'} \bar{S}_{n-1}, \mathcal{E}_\lambda) = \bigoplus_{w \in W(G')} H^q(\mathfrak{g}', K', \text{Ind}_{B'(\mathbb{A}_{\mathcal{K}})}^{G'(\mathbb{A}_{\mathcal{K}})}[\tilde{\tau}^w] \otimes E_\lambda) \oplus \bigoplus_{\substack{\chi \neq \tilde{\tau}^w \\ \forall w \in W(G')}} H^q(\mathfrak{g}', K', \text{Ind}_{B'(\mathbb{A}_{\mathcal{K}})}^{G'(\mathbb{A}_{\mathcal{K}})}[\chi] \otimes E_\lambda),$$

and there is no weak $G'(\mathbb{A}_f)$ -intertwining between the two sums. Hence, for degree $q = b_{n-1}$ we obtain

$$(5.1) \quad H^{b_{n-1}}(\partial_{B'} \bar{S}_{n-1}, \mathcal{E}_\lambda) = \bigoplus_{w \in W(G')} \text{Ind}_{B'(\mathbb{A}_f)}^{G'(\mathbb{A}_f)}[\tilde{\tau}^w] \oplus \bigoplus_{\substack{\chi \neq \tilde{\tau}^w \\ \forall w \in W(G')}} H^{b_{n-1}}(\mathfrak{g}', K', \text{Ind}_{B'(\mathbb{A}_{\mathcal{K}})}^{G'(\mathbb{A}_{\mathcal{K}})}[\chi] \otimes E_\lambda),$$

and the summands in the first sum are all irreducible.

Proposition 5.2. *For all $\sigma \in \text{Aut}(\mathbb{C})$, the following diagram of $G'(\mathbb{A}_f)$ -equivariant homomorphisms commutes:*

$$\begin{array}{ccccccc} H^q(\mathfrak{g}', K', \Pi' \otimes E_\lambda) & \hookrightarrow & H^q(\bar{S}_{n-1}, \mathcal{E}_\lambda) & \xrightarrow{\text{res}^q} & H^q(\partial \bar{S}_{n-1}, \mathcal{E}_\lambda) & \xrightarrow{\text{res}_{B'}^q} & H^q(\partial_{B'} \bar{S}_{n-1}, \mathcal{E}_\lambda) \\ \downarrow \sigma^* & & \downarrow \sigma^* & & \downarrow \sigma^* & & \downarrow \sigma^* \\ H^q(\mathfrak{g}', K', {}^\sigma \Pi' \otimes {}^\sigma E_\lambda) & \hookrightarrow & H^q(\bar{S}_{n-1}, {}^\sigma \mathcal{E}_\lambda) & \xrightarrow{\text{res}^q} & H^q(\partial \bar{S}_{n-1}, {}^\sigma \mathcal{E}_\lambda) & \xrightarrow{\text{res}_{B'}^q} & H^q(\partial_{B'} \bar{S}_{n-1}, {}^\sigma \mathcal{E}_\lambda) \end{array}$$

Furthermore, there are sections $s_{B'}$ to the composition $r_{B'}$ of the morphisms in the rows of the diagram,

$$s_{B'} : H^{b_{n-1}}(\partial_{B'} \bar{S}_{n-1}, {}^\sigma \mathcal{E}_\lambda) \longrightarrow H^{b_{n-1}}(\mathfrak{g}', K', {}^\sigma \Pi' \otimes {}^\sigma E_\lambda).$$

Proof. We only prove the last assertion. Recall the archimedean vectors $\xi_{\Pi'_\infty, \underline{j}, \beta} \in W(\Pi'_\infty)$ from Sect. 3.3. Then, for every non-zero $\xi_f \in W(\Pi'_f)$, we let

$$E_{\Pi', \underline{j}, \beta} := W^{-1}(\xi_{\Pi'_\infty, \underline{j}, \beta} \otimes \xi_f) \in \Pi'$$

be the corresponding Eisenstein series. Hence,

$$\sum_{\underline{j}, \beta} \left(X_{\underline{j}}'^* \otimes E_{\Pi', \underline{j}, \beta} \otimes e'_\beta \right) \in H^{b_{n-1}}(\mathfrak{g}', K', \Pi' \otimes E_\lambda)$$

is a non-zero cohomology class by the choice of $\xi_{\Pi'_\infty, \underline{j}, \beta}$. Via the inclusion

$$H^{b_{n-1}}(\mathfrak{g}', K', \Pi' \otimes E_\lambda) \hookrightarrow H^{b_{n-1}}(\bar{S}_{n-1}, \mathcal{E}_\lambda)$$

it is mapped onto

$$\sum_{\underline{j}, \beta} \left(E_{\Pi', \underline{j}, \beta} \otimes e'_\beta \right) dx_{\underline{j}} \in H^{b_{n-1}}(\bar{S}_{n-1}, \mathcal{E}_\lambda)$$

and finally, when restricting to the Borel-stratum, we obtain

$$\sum_{\underline{j}, \beta} \left(E_{\Pi', \underline{j}, \beta}|_{B'} \otimes e'_\beta \right) dx_{\underline{j}} \in H^{b_{n-1}}(\partial_{B'} \bar{S}_{n-1}, \mathcal{E}_\lambda).$$

Here, $E_{\Pi', \underline{j}, \beta}|_{B'}$ is the constant term of the Eisenstein series $E_{\Pi', \underline{j}, \beta}$ with respect to the Borel subgroup B' , see [45] Satz 1.10. According to [38] Prop. II.7 this constant term can be written as

$$E_{\Pi', \underline{j}, \beta}|_{B'} = \sum_{w \in W(G')} M(w, \tilde{\tau}) f_{\Pi', \underline{j}, \beta},$$

where $f_{\Pi', \underline{j}, \beta}$ is a K' -finite section in $\text{Ind}_{B'(\mathbb{A}_K)}^{G'(\mathbb{A}_K)}[\tilde{\tau}]$, used to define the Eisenstein series $E_{\Pi', \underline{j}, \beta}$ and $M(w, \tilde{\tau})$ is the corresponding intertwining operator,

$$M(w, \tilde{\tau}) : \text{Ind}_{B'(\mathbb{A}_K)}^{G'(\mathbb{A}_K)}[\tilde{\tau}] \rightarrow \text{Ind}_{B'(\mathbb{A}_K)}^{G'(\mathbb{A}_K)}[\tilde{\tau}^w]$$

cf. [38] II.6. As a consequence, the Eisenstein cohomology class, when restricted to the Borel-stratum, yields the class

$$\sum_{w \in W(G')} \sum_{\underline{j}, \beta} \left(M(w, \tilde{\tau}) f_{\Pi', \underline{j}, \beta} \otimes e'_\beta \right) dx_{\underline{j}},$$

which, via the isomorphism (5.1), lies in $\bigoplus_{w \in W(G')} \text{Ind}_{B'(\mathbb{A}_f)}^{G'(\mathbb{A}_f)}[\tilde{\tau}^w]$. Here we observe that there is no weak intertwining of the two sums in (5.1). This class is non-zero, since all Eisenstein series $E_{\Pi', \underline{j}, \beta}$ are holomorphic at their point of evaluation $s = 0$, cf. the proof of Thm. 4.11 in [45]. Hence, since all representations $\text{Ind}_{B'(\mathbb{A}_f)}^{G'(\mathbb{A}_f)}[\tilde{\tau}^w]$ are irreducible, the various summands of the image of $r_{B'}$ are either trivial or the whole target space. The assertion follows now from the irreducibility of the source $\Pi'_f = H^{b_{n-1}}(\mathfrak{g}', K', \Pi' \otimes E_\lambda)$. \square

5.2. A rationality result. Recall Prop. 2.5. It is clear that $H^{b_{n-1}}(\partial_{B'} \bar{S}_{n-1}, \mathcal{E}_\lambda)$ inherits from E_λ a natural $\mathbb{Q}(E_\lambda)$ -structure. By Prop. 2.4 and the temperedness of Π'_∞ , there is the equality of number fields $\mathbb{Q}(E_\lambda) = \mathbb{Q}(\Pi'_\infty)$. Hence, by strong multiplicity one for isobaric automorphic representations, $\mathbb{Q}(E_\lambda) \subseteq \mathbb{Q}(\Pi'_f)$. Next, observe that by Prop. 2.5, $\Pi'_f = H^{b_{n-1}}(\mathfrak{g}', K', \Pi' \otimes E_\lambda)$ carries a natural $\mathbb{Q}(\Pi'_f)$ -structure. We deduce the following corollary.

Corollary 5.3. *The morphism $r_{B'}$ is $\mathbb{Q}(\Pi'_f)$ -rational, i.e., it maps the natural $\mathbb{Q}(\Pi'_f)$ -structure of the cohomology space $H^{b_{n-1}}(\mathfrak{g}', K', \Pi' \otimes E_\lambda)$ to the natural $\mathbb{Q}(\Pi'_f)$ -structure of boundary cohomology $H^{b_{n-1}}(\partial_{B'} \bar{S}_{n-1}, \mathcal{E}_\lambda)$. The same holds true for the section $s_{B'}$.*

Proof. This is clear by the commutativity of the diagram in Prop. 5.2. \square

Remark 5.4. We would like to point out that for regular coefficients E_λ , Thm. 7.23 in Grobner–Raghuram [16] should provide an alternative approach to Cor. 5.3.

Recall from Section 3.4 that the rational structure on Π'_f is given by the natural rational structure on the induced representation. Let $F \mapsto F|_{B'}$ denote the constant term on functions (Eisenstein series) $F \in \Pi'$, and let $F|_{B'}(\tilde{\tau})$ denote its $\tilde{\tau}$ -component with respect to the action of the maximal torus in B' . It follows from the proof of Proposition 5.2 that

Lemma 5.5. *Let $F = F_\infty \otimes F_f \in \Pi'$ be a function with F_∞ in the K' -type contributing to the cohomology space $H^{b_{n-1}}(\mathfrak{g}', K', \Pi'_\infty \otimes E_\lambda)$. Let $L \supseteq \mathbb{Q}(\Pi'_f)$ be a field extension. Then F defines an L -rational cohomology class $[F]$ if and only if F_f , viewed as an element of $\text{Ind}_{B'(\mathbb{A}_f)}^{G'(\mathbb{A}_f)}[\tilde{\tau}_f]$, takes values in L , which is furthermore equivalent to $F|_{B'}(\tilde{\tau})$, viewed as an element of $\text{Ind}_{B'(\mathbb{A}_f)}^{G'(\mathbb{A}_f)}[\tilde{\tau}_f]$, taking values in L . Moreover, for all $\sigma \in \text{Aut}(\mathbb{C})$, $\sigma[F] = {}^\sigma F_\infty \otimes \sigma(F_f)$ (the notation having the obvious meaning).*

5.3. Whittaker coefficients. Let $f = \otimes'_w f_w$ be a K' -finite, decomposable section in $\text{Ind}_{B'(\mathbb{A}_K)}^{G'(\mathbb{A}_K)}[\tilde{\tau}]$. We let $E(f) = E(f, 0)$ be the attached Eisenstein series, evaluated at 0, where it is always holomorphic. We will consider its global ψ -Whittaker functional (ψ -Fourier coefficient):

$$E_\psi(f)(g) := \int_{U'(\mathcal{K}) \backslash U'(\mathbb{A}_K)} E(f)(u'g)\psi^{-1}(u')du' \in W(\Pi').$$

To each “function” $\varphi \in \tilde{\tau}$ (which is of course simply a constant), we can associate a K' -finite, decomposable section $f_\varphi = \otimes'_w f_{\varphi,w} \in \text{Ind}_{B'(\mathbb{A}_K)}^{G'(\mathbb{A}_K)}[\tilde{\tau}]$ as in [44] 6.3. Recall the choice of an element $w_0 \in G'(\mathcal{K})$, representing the longest element in the Weyl group of G'/\mathcal{K} . Then we obtain the following result

Proposition 5.6. *The value at the identity $id \in G'(\mathbb{A}_K)$ of the ψ -Whittaker functional $E_\psi(f_\varphi)$ equals*

$$E_\psi(f_\varphi)(id) = \prod_{w \in S(\Pi')} W_w(id_w) \cdot \prod_{\substack{w \notin S(\Pi') \\ 1 \leq i < j \leq n-1}} L(1, BC(\chi_i)_w BC(\chi_j^{-1})_w)^{-1},$$

where

$$W_w(id_w) = \int_{U'(\mathcal{K}_w)} f_{\varphi,w}(w_0^{-1}n'_w)\psi_w^{-1}(n'_w)dn'_w.$$

In particular, $E_\psi(f_\varphi)(id)$ has an Euler factorization.

Proof. This is [44] Thm. 7.1.2. One simply observes that Shahidi’s functional λ_w^0 (cf. [44], p. 122) gives the complex number $f_{\varphi,w}(w_0^{-1}n'_w)$, viewed as an element in $\tilde{\tau}$. \square

We recall the \mathbb{Q} -rational (and hence by our assumption also $\mathbb{Q}(E_\lambda)$ -rational) basis vectors $X_{\underline{j}}'^*$, resp. the $\mathbb{Q}(E_\lambda)$ -basis $\{e'_\beta\}$ of E_λ from Sect. 3.3. Moreover, we assume from now on that $\xi_{\Pi'_\infty, \underline{j}, \beta}$ was chosen in a way such that the generator $[\Pi'_\infty]$ is $\mathbb{Q}(E_\lambda)$ -rational in the sense of Sect. 2.7.

We now choose a section f_φ in such a way that $f_{\varphi, \infty}$ is one of the vectors $W^{-1}(\xi_{\Pi'_\infty, \underline{j}, \beta})$ appearing in the definition of $[\Pi'_\infty]$ (or rather its transfer by W^{-1}) and that the non-archimedean part of f_φ lies in the given $\mathbb{Q}(\Pi'_f)$ -structure of Π'_f . Then, the evaluated Eisenstein series $E(f_\varphi)$ appears as a factor in the tensor product $\sum_{\underline{j}, \beta} (X_{\underline{j}}'^* \otimes E_{\Pi', \underline{j}, \beta} \otimes e'_\beta)$, which defines a $\mathbb{Q}(\Pi'_f)$ -rational Eisenstein cohomology class.

Corollary 5.7. *There is a constant $\Omega(\Pi'_\infty) \in \mathbb{C}^\times$, depending only on Π'_∞ and on the choice of basis vectors $\xi_{\Pi'_\infty, j, \beta}$, such that*

$$p(\Pi') \sim_{\mathbb{Q}(\Pi'_f)} \Omega(\Pi'_\infty) \cdot \prod_{1 \leq i < j \leq n-1} L(1, BC(\chi_i \cdot \chi_j^{-1})_f).$$

Proof. It follows from Corollary 5.3, 5.5, and the definitions that the period $p(\Pi')$ can be represented by the ratio between the value at the identity of the $\tilde{\tau}$ -component of the constant term and the value at the identity of the ψ -Whittaker functional $E_\psi(f_\varphi)$, if the latter doesn't vanish. We have chosen f_φ so that $E(f_\varphi)$ defines a $\mathbb{Q}(\Pi'_f)$ -rational class, which means that $p(\Pi')$ is the inverse of $E_\psi(f_\varphi)(id)$, which by Proposition 5.6 equals

$$W_\infty(id_\infty) \cdot \prod_{\substack{w \in S(\Pi') \setminus \{\infty\} \\ 1 \leq i < j \leq n-1}} W_w(id_w) \cdot L(1, BC(\chi_i)_w BC(\chi_j^{-1})_w) \cdot \prod_{1 \leq i < j \leq n-1} L(1, BC(\chi_i)_f BC(\chi_j^{-1})_f)^{-1}.$$

We take $\Omega(\Pi'_\infty) = W_\infty(id_\infty)^{-1}$. Since the local L -factors belong to $\mathbb{Q}(\Pi'_f)$ and transform under the Galois group along with Π' , we will finish the proof if we can show that $f_{\varphi, w}$ can be chosen so that the integral defining $W_w(id_w)$ is a rational constant. Now, as in Lemma 5.5, the $\mathbb{Q}(\Pi'_w)$ -rational structure on the principal series representation Π'_w is given by that on the boundary cohomology, and coincides with the one defined (adelically) in 2.6, namely the space of functions in the principal series whose restrictions to a fixed maximal compact subgroup K'_w take values in $\mathbb{Q}(\Pi'_w)$. We take $U_1 \subset U'(\mathcal{K}_w)$ to be a subgroup of the kernel of ψ_w . Possibly shrinking U_1 further, we can assume that $f_{\varphi, w}(w_0 n'_w)$ is supported in U_1 as a function of n'_w and takes value 1 there. Then the integral is just a volume factor and belongs to \mathbb{Q} . \square

6. PERIOD RELATIONS - THE MAIN RESULTS

In this section we always assume Π and Π' to be obtained by base change from unitary groups of all signatures; in other words, that they both satisfy Hypothesis 4.19.

6.1. Critical values of tensor products when one of the representations is of abelian type. Let Π' be the abelian automorphic representation of 2.5, attached to the $(n-1)$ -tuple $\chi = (\chi_1, \dots, \chi_{n-1})$ and the auxiliary character γ ; we write $\Pi' = \Pi'(\chi, \gamma)$. Recall that $\chi_{j, \infty}(e^{i\theta}) = e^{ik_j \theta}$ and $\gamma_\infty(z) = (z/\bar{z})^t$ for $k_j \in \mathbb{Z}$ and $t \in \frac{n}{2} + \mathbb{Z}$. If n is odd, we henceforth assume $t = \frac{1}{2}$, to apply the formulas of [22].

Define $\gamma^+(x)$ as in Rem. 2.6, i.e., $\gamma^+(x) = \gamma(x) \cdot \|x\|^{\frac{1}{2}}$ if n is odd and $\gamma^+(x) = \gamma(x)$ if n is even. Then, if n is odd:

$$(6.1) \quad \begin{aligned} L(s, R(M(\Pi) \otimes M(\Pi'))) &= \prod_{j=1}^{n-1} L(s + \frac{3-2n}{2}, \Pi \otimes BC(\chi_j) \cdot \gamma) \\ &= \prod_{j=1}^{n-1} L(s - \frac{n-1}{2}, M(\Pi) \otimes M(BC(\chi_j) \cdot \gamma^+)). \end{aligned}$$

Moreover, if n is odd, $\gamma_\infty^+(z) = z$ and thus, γ^+ is of type η_ℓ with $\ell = -1$, in the notation of [21] (2.9.7) and (especially) Theorem 4.3 of [22], see p. 177. If n is even, then $\gamma_\infty^+(z) = 1$, so γ^+ is of type η_ℓ with $\ell = 0$. In what follows, we let $\alpha = \gamma^+$ and so the restriction $\alpha_0 = \varepsilon_{\mathcal{K}}^n$

for all n . Thus, with this choice of α , the term $(2\pi i)^{\kappa r(M')} \mathcal{G}(\alpha_{0,f}^{-1})^{r(M')}$ in Thm. 4.31 is just $(2\pi i)^{\kappa r(M')} \cdot \mathcal{G}(\varepsilon_{\mathcal{K},f}^n)^{r(M')}$. Let $s_0 = n - 1 + m$, where m is as in 4.7 above. We apply Thm. 4.31 to the critical values of $L(s, R(M(\Pi) \otimes M(\Pi'))) = L(s, R(M(\Pi^\vee) \otimes M(\Pi'^c)))$. The results are as follows. If n is even, $\alpha = \gamma^+$ is the trivial character and we have

$$\begin{aligned}
(6.2) \quad L(n-1+m, R(M(\Pi) \otimes M(\Pi'))) &= \prod_{j=1}^{n-1} L(\frac{n}{2} + m, M(\Pi^\vee) \otimes M(BC(\chi_j)^c)) \\
&\sim_{\mathcal{K}E(\Pi)E(\Pi')} \prod_{j=1}^{n-1} (2\pi i)^{(\frac{n}{2}+m)n - \frac{n(n-1)}{2}} \cdot \mathcal{G}(\varepsilon_{\mathcal{K},f})^{\frac{n}{2}} \cdot P^{(r(M_j^+))}(\Pi) G(\Pi_j^{+,c})^{n-2r(M_j^+)} \\
&\sim_{\mathcal{K}E(\Pi)E(\Pi')} (2\pi i)^{(n-1)(mn) + \frac{n(n-1)}{2}} \mathcal{G}(\varepsilon_{\mathcal{K},f})^{\frac{n(n-1)}{2}} \prod_{j=1}^{n-1} P^{(r(M_j^+))}(\Pi) G(\Pi_j^{+,c})^{n-2r(M_j^+)}, \\
&\sim_{\mathcal{K}E(\Pi)E(\Pi')} (2\pi i)^{(n-1)(mn) + \frac{n(n-1)}{2}} \mathcal{G}(\varepsilon_{\mathcal{K},f})^{\frac{n(n-1)}{2}} \prod_{j=1}^{n-1} P^{(j)}(\Pi) G(\Pi_j^{+,c})^{n-2j}
\end{aligned}$$

with $\Pi_j^{+,c} = (BC(\chi_j)\gamma^+)^c = BC(\chi_j)^{-1} = \Pi_j'^c$ and $M_j^+ = M(\Pi_j^+) = M_j'$. In order to obtain the last line, note that the character χ_j has infinity type $(z/\bar{z})^{k_j}$ which in the conventions of 4.2 means that the parameter is $-k_j$. Then the set $T(M, M_j')$ is the set of t such that $2k_j \geq n - 2p_t^c$. On the other hand, we are given that $2k_j \in [n - 2q_{j+1}, 2p_j - n]$, i.e., j is the largest integer such that $2k_j \geq n - 2q_{j+1} = n - 2p_{n-j}^c$. Thus, $s(M_j^+) = n - j$ by (4.16) and so $r(M_j^+) = r(M_j') = j$. If n is odd, then $\alpha = \gamma^+$ and we have

$$\begin{aligned}
(6.3) \quad L(n-1+m, R(M(\Pi) \otimes M(\Pi'))) &= \prod_{j=1}^{n-1} L(\frac{n-1}{2} + m, M(\Pi^\vee) \otimes M((BC(\chi_j)\gamma^+)^c)) \\
&\sim_{\mathcal{K}E(\Pi)E(\Pi')} \prod_{j=1}^{n-1} (2\pi i)^{mn+r(M_j^+)} \mathcal{G}(\varepsilon_{\mathcal{K},f})^{\lceil \frac{n}{2} \rceil + r(M_j^+)} P^{(r(M_j^+))}(\Pi) G(\Pi_j^{+,c})^{n-2r(M_j^+)} \\
&\sim_{\mathcal{K}E(\Pi)E(\Pi')} (2\pi i)^{(n-1)(mn) + \frac{n(n-1)}{2}} \mathcal{G}(\varepsilon_{\mathcal{K},f})^{\lceil \frac{n}{2} \rceil (n-1) + \frac{n(n-1)}{2}} \prod_{j=1}^{n-1} P^{(j)}(\Pi) G(\Pi_j^{+,c})^{n-2j}
\end{aligned}$$

Here $\Pi_j^{+,c} = (BC(\chi_j)\gamma^+)^c = BC(\chi_j)^{-1}(\gamma^+)^c$. The contribution of γ^+ disappears because the product of the periods of γ^+ to the $n - 2j$ is just 1. In particular, the expressions (6.2) and (6.3) are identical (up to the power $\mathcal{G}(\varepsilon_{\mathcal{K},f})^{\frac{n(n-1)}{2}}$ of the Gauß-sum, which we may absorb into $\mathcal{K}E(\Pi)E(\Pi')$).

6.2. Whittaker periods and Petersson norms. In this section we combine the results of the previous sections to express the invariants $p(\Pi)$ in terms of the $P^{(s)}(\Pi)$. Notation is as above; in particular, Π is cohomological and obtained by base change from unitary groups and $\Pi' = \Pi'(\chi, \gamma)$.

Proposition 6.4. *Suppose Π satisfies Hypothesis 4.19. Let $s_0 = n - 1 + m$, $m \geq 0$, be a critical value of $L(s, R(M(\Pi) \otimes M(\Pi')))$ satisfying $L(n - 1 + m, R(M(\Pi) \otimes M(\Pi'))) \neq 0$.*

Then there are constants $p(m, \Pi_\infty, \Pi'_\infty) \in \mathbb{C}^\times$ and $\Omega(\Pi'_\infty) \in \mathbb{C}^\times$ such that

$$(6.5) \quad p(m, \Pi_\infty, \Pi'_\infty) \Omega(\Pi'_\infty) p(\Pi) \prod_{1 \leq i < j \leq n-1} L(1, BC(\chi_i \chi_j^{-1})_f) \sim_{\mathcal{KE}(\Pi)E(\Pi')} \\ (2\pi i)^{(n-1)(mn) + \frac{n(n-1)}{2}} \mathcal{G}(\varepsilon_{\mathcal{K}, f})^{\lceil \frac{n}{2} \rceil (n-1)} \prod_{j=1}^{n-1} P^{(j)}(\Pi) G(\Pi'^c)^{n-2j}.$$

A critical point $s_0 = n - 1 + m$, $m \geq 0$, giving $L(n - 1 + m, R(M(\Pi) \otimes M(\Pi'))) \neq 0$ exists in particular, if the inequalities separating the archimedean parameters of Π and Π' are strict:

$$\mu_1 > -\lambda_{n-1} > \mu_2 > -\lambda_{n-2} > \dots > -\lambda_1 > \mu_n.$$

Proof. This is obtained by comparing the expressions of 3.9, 5.7, 6.2, and 6.3 for the non-vanishing value of the L -function, and bearing in mind that, by 3.10, the term $\mathcal{G}(\omega_{\Pi'_{f,0}})$ in 3.9 is trivial. The last assertion follows from the description of 4.7: It implies there is a critical value s_0 strictly to the right of the center of symmetry. Since Π is cuspidal, it is a theorem of Shahidi and Jacquet-Shalika that the value $L(s_0, R(M(\Pi) \otimes M(\Pi'))) \neq 0$. \square

Now, let $M'_{ij} := M(BC(\chi_i \chi_j^{-1}))$ with $i < j$. Then, the parameter of M'_{ij} is $2(k_j - k_i) < 0$. So $r(M'_{ij}) = 0$ and we get

$$L(1, M'_{ij}) \sim (2\pi i) G(BC(\chi_i \chi_j^{-1})^c) = (2\pi i) G(\Pi'_i{}^c) G(\Pi'_j{}^c)^{-1}.$$

Then

$$\prod_{1 \leq i < j \leq n-1} L(1, M'_{ij}) \sim (2\pi i)^{\frac{(n-1)(n-2)}{2}} \prod_{j=1}^{n-1} G(\Pi'_j{}^c)^{n-2j}.$$

Putting this together we find that, assuming the hypotheses of the preceding proposition are verified, we have

$$p(m, \Pi_\infty, \Pi'_\infty) \Omega(\Pi'_\infty) p(\Pi) (2\pi i)^{\frac{(n-1)(n-2)}{2}} \prod_{j=1}^{n-1} G(\Pi'_j{}^c)^{n-2j} \sim_{\mathcal{KE}(\Pi)E(\Pi')} \\ (2\pi i)^{(n-1)(mn) + \frac{n(n-1)}{2}} \mathcal{G}(\varepsilon_{\mathcal{K}, f})^{\lceil \frac{n}{2} \rceil (n-1)} \prod_{j=1}^{n-1} P^{(j)}(\Pi) \prod_{j=1}^{n-1} G(\Pi'_j{}^c)^{n-2j}$$

or more simply

$$(6.6) \quad p(m, \Pi_\infty, \Pi'_\infty) \Omega(\Pi'_\infty) p(\Pi) \sim_{\mathcal{KE}(\Pi)E(\Pi')} (2\pi i)^{(nm+1)(n-1)} \prod_{j=1}^{n-1} P^{(j)}(\Pi).$$

Now, we can state our first main theorem:

Theorem 6.7. *Let μ be the parameter of Π , and suppose $\mu_i - \mu_{i+1} \geq 2$ for all i , so that there is a parameter λ for G' and an integer $m > 0$ such that $s_0 = n - 1 + m$ is a critical value of the tensor product L -function $L(s, R(M(\Pi) \otimes M(\Pi')))$ (in the motivic normalization) for*

any Π' with parameter λ . Suppose Π satisfies Hypothesis 4.19. Then there is a non-zero constant $Z(\Pi_\infty)$ which depends only on the local representation Π_∞ and such that

$$p(\Pi) \sim_{\mathcal{KE}(\Pi)} Z(\Pi_\infty) \prod_{j=1}^{n-1} P^{(j)}(\Pi).$$

Proof. The hypothesis on μ implies that there exist Π' of abelian type such that the comparison of (6.6) is valid. By letting Π' vary among abelian type representations with parameter λ , we can remove the $E(\Pi')$ from the equivalence relation. Initially one obtains the relation

$$(6.8) \quad p(\Pi) \sim_{\mathcal{KE}(\Pi)} Z(m, \Pi_\infty, \Pi'_\infty) \prod_{j=1}^{n-1} P^{(j)}(\Pi),$$

where $Z(m, \Pi_\infty, \Pi'_\infty) = [\Omega(\Pi'_\infty)p(m, \Pi_\infty, \Pi'_\infty)]^{-1} \cdot (2\pi i)^{(nm+1)(n-1)}$. But since the left-hand side of 6.8 is independent of m and Π'_∞ , so is the right-hand side. \square

Remark 6.9. Two substitutes are possible for Hypothesis 4.19 when it is not satisfied. As noted above, there is only an obstruction if n is even. Under the regularity hypothesis on the parameter μ , it can be shown that the standard L -function of Π can be realized, up to an abelian twist, as the L -function of a holomorphic automorphic representation of $U(n+1-j, j)$ for any j ; this was done in most cases using the theta correspondence in [23]. The Petersson norm of an arithmetically normalized form in this representation can then be used in place of the missing $P^{(j)}(\Pi)$.

Even in the absence of the regularity hypothesis, one can use quadratic base change, as in [49], to define versions of the missing $P^{(j)}(\Pi)$. These are only well-defined up to square roots of elements in the coefficient field, so are less precise than the ones defined motivically.

6.3. General tensor products. Combining the comparison in Thm. 6.7 with Thm. 1.3, we obtain the following general result, when Π_∞ and Π'_∞ are both sufficiently regular.

Theorem 6.10. *Let $\Pi = BC(\pi)$ and $\Pi' = BC(\pi')$ be cuspidal automorphic representations of $G(\mathbb{A}_K)$ and $G'(\mathbb{A}_K)$ which are cohomological with respect to E_μ and E_λ , respectively. Suppose*

- (1) $\mu_i - \mu_{i+1} \geq 2$ for all i and $\lambda_j - \lambda_{j+1} \geq 2$ for all j .
- (2) Both Π and Π' satisfy Hypothesis 4.19.

Then for every critical point $s_0 = \frac{1}{2} + m$ of $L(s, \Pi \times \Pi')$ with $m \geq 0$,

$$L(\tfrac{1}{2} + m, \Pi_f \times \Pi'_f) \sim_{\mathcal{KE}(\Pi)E(\Pi')} p(m, \Pi_\infty, \Pi'_\infty) Z(\Pi_\infty) Z(\Pi'_\infty) \prod_{j=1}^{n-1} P^{(j)}(\Pi) \prod_{k=1}^{n-2} P^{(k)}(\Pi').$$

Equivalently, for every critical point $s_0 = n - 1 + m$ of $L(s, R(M(\Pi) \otimes M(\Pi')))$ with $m \geq 0$,

$$L(n - 1 + m, R(M(\Pi) \otimes M(\Pi'))) \sim_{\mathcal{KE}(\Pi)E(\Pi')} p(m, \Pi_\infty, \Pi'_\infty) Z(\Pi_\infty) Z(\Pi'_\infty) \prod_{j=1}^{n-1} P^{(j)}(\Pi) \prod_{k=1}^{n-2} P^{(k)}(\Pi').$$

6.4. The archimedean constant. The constants $\Omega(\Pi'_\infty)$, $p(m, \Pi_\infty, \Pi'_\infty)$, and $p(\Pi)$ all depend on choices of test vectors in the induced and Whittaker models of Π_∞ and Π'_∞ . We have decided to choose vectors rational over \mathcal{K} , but we could also choose test vectors with the property that $W_\infty(\Pi') = 1$. Since this choice depends only on Π'_∞ , it makes sense for cuspidal as well as Eisenstein representations. This provides a unique normalization of $p(\Pi)$ and thus a unique value for $p(m, \Pi_\infty, \Pi'_\infty)$ (as does the choice of rational vectors).

In [35], Jie Lin proves a period relation for Π and Π' obtained by automorphic induction from Hecke characters. Using this relation, she deduces the following refinement of Theorem 6.10.

Theorem 6.11 (Lin). *Under the hypotheses of Theorem 6.10, we have the following relations:*

$$L\left(\frac{1}{2} + m, \Pi_f \times \Pi'_f\right) \sim_{\mathcal{K}E(\Pi)E(\Pi')} (2\pi i)^{(m+\frac{1}{2})n(n-1)} \prod_{j=1}^{n-1} P^{(j)}(\Pi) \prod_{k=1}^{n-2} P^{(k)}(\Pi').$$

or equivalently,

$$L\left(n-1+m, R(M(\Pi) \otimes M(\Pi'))\right) \sim_{\mathcal{K}E(\Pi)E(\Pi')} (2\pi i)^{(m+\frac{1}{2})n(n-1)} \prod_{j=1}^{n-1} P^{(j)}(\Pi) \prod_{k=1}^{n-2} P^{(k)}(\Pi'),$$

provided $m > 0$.

If $m = 0$ her result remains true under a certain global non-vanishing hypothesis. In view of formula (4.12), this is exactly what is predicted by Deligne's conjecture, assuming the validity of Optimistic Comparison 4.32.

6.5. Whittaker periods and adjoint L -values. The results of this section, in contrast to Thm. 6.7, are unconditional, but for the moment they only apply under slightly restrictive hypotheses. This is because they are based on Wei Zhang's version [53] of the Ichino-Ikeda-Neal Harris conjecture for automorphic forms on unitary groups, see [20], and for the moment neither the local nor global properties of the relative trace formula are known in sufficient generality to allow a complete comparison. The simplifying hypotheses in [53] are analogous to those used in earlier applications of the Arthur-Selberg trace formula, and the history of the latter gives us reason to be optimistic that the restrictions will soon be unnecessary.

For ease of reference, the simplifying hypotheses are listed separately.

6.5.1. Simplifying hypotheses. The unitary groups H and H' are as in the earlier sections. We let π and π' be unitary cuspidal automorphic representations of $H(\mathbb{A})$, resp. $H'(\mathbb{A})$, which occur with multiplicity one in the discrete spectrum of H , resp. H' . Say a prime p is *split* if it splits in the quadratic extension \mathcal{K} . Since we want to use Wei Zhang's results, we have to assume

- Hypothesis 6.12.**
- (a) There exists some split prime p such that π_p and π'_p are both supercuspidal.
 - (b) Every prime $p < M$ is split, where M is the "algorithmically computable" constant that arises in [13] in the transfer to characteristic zero of the Jacquet-Rallis fundamental lemma (see below).
 - (c) If p is not split, then π_p and π'_p are both unramified.

Remark 6.13. (a) Let $\Pi = BC(\pi)$ and $\Pi' = BC(\pi')$ denote the base change representations of π and π' , to $G(\mathbb{A}_{\mathcal{K}})$ and $G'(\mathbb{A}_{\mathcal{K}})$, respectively. Hypothesis (a) above implies that Π and Π' are both cuspidal automorphic representations. Moreover, by Thm. 2.2, Π and Π' are both of cohomological type. Hence, the Asai L -functions $L(s, \Pi, As^{(-1)^{n-1}})$ and $L(s, \Pi', As^{(-1)^{n-2}})$ have a pole at $s = 1$, by [25] or [39] Cor. 2.5.9 and (2.5.12). As a consequence, Wei Zhang's additional set of assumptions **RH(I)** (cf. [53], p. 544) is satisfied in our case. Moreover, Zhang's second set of assumptions **RH(II)** (cf. [53], pp. 544–545) is fulfilled by the work of Beuzart-Plessis, [2]. (We should, however, mention that the latter paper depends on the (expected) properties of local L -packets for unitary groups. See [2] Sect. 18.1 for the precise expectations.)

(b) In [53] it is also assumed that π and π' are locally tempered at all places. Due to Caraiani [5], Thm. 1.2, $\Pi = BC(\pi)$ and $\Pi' = BC(\pi')$ are tempered at all places (under hypothesis (a) it was also proved in the book of Harris and Taylor). We expect that by [31] and the forthcoming work of Kaletha, Minguez, Shin and White, it should automatically follow that π and π' are also tempered at all places, whence we did not assume this explicitly. However, the extremely careful reader may restrict himself/herself in what follows to the case where π and π' are locally tempered at all primes p .

(c) Zhiwei Yun proves the Jacquet-Rallis fundamental lemma in [51] for local fields of characteristic $> n$. In her appendix to Yun's paper, Julia Gordon derives the Jacquet-Rallis fundamental lemma for a local field K of characteristic 0, but at present her methods (from motivic integration) require her to assume that the residue characteristic of K is greater than an unspecified positive constant M . This is the M that appears in (b) above; it is effectively computable but no one has carried out the computation. For split primes the Jacquet-Rallis fundamental lemma is vacuous. The requirement that a prime p be split in the results of [53] can be removed whenever the Jacquet-Rallis fundamental lemma is known in residue characteristic p .

Since $H(\mathbb{R})$ and $H'(\mathbb{R})$ are both assumed compact we are in case (2) of [53], Theorem 1.2. We define

$$\mathcal{L}(\pi, \pi') := \Delta_n \cdot \frac{L(\frac{1}{2}, \Pi \times \Pi')}{L(1, \pi, Ad)L(1, \pi', Ad)},$$

where Δ_n is the L -function of the *Gross motive*:

$$\Delta_n = \prod_{i=1}^n L(i, \varepsilon_{\mathcal{K}, f}^i).$$

For $\phi \in \pi$ and $\phi' \in \pi'$, we let

$$P(\phi, \phi') := \int_{H'(\mathbb{Q}) \backslash H'(\mathbb{A})} \phi(h') \phi'(h') dh';$$

$$\langle \phi, \phi \rangle := \int_{H(\mathbb{Q}) \backslash H(\mathbb{A})} \phi(h) \bar{\phi}(h) dh.$$

We use Tamagawa measures for dh and dh' . Here is Wei Zhang's theorem.

Theorem 6.14. ([53]) *Assume π , π' , and \mathcal{K} satisfy hypothesis 6.12. Then there is a non-zero constant $c(\pi_{\infty}, \pi'_{\infty})$, depending only on the archimedean components of π and π' , such*

that, for every factorizable $\phi \in \pi$ and $\phi' \in \pi'$,

$$2^{-2}c(\pi_\infty, \pi'_\infty)Z_{loc}(\phi, \phi')\mathcal{L}(\pi, \pi') = \frac{|P(\phi, \phi')|^2}{\langle \phi, \phi \rangle \langle \phi', \phi' \rangle}.$$

Here $Z_{loc} = \prod_{v \in S} Z_v$ is a product of normalized local integrals of matrix coefficients; Z_v is denoted

$$\frac{\alpha_v^{\natural}(\Phi_v, \Phi_v)}{(\Phi_v, \Phi_v)_v}$$

in [53], with $\Phi_v = \phi_v \otimes \phi'_v$ (Zhang uses ϕ_v for this tensor product).

Remark 6.15. The theorem in [53] is stated under a more general version of hypothesis (c). With a bit more work we could derive the consequences below in this more general situation; however, hypothesis (c) is destined to disappear in the short term, so this seems unnecessary.

As in (4.1.2) of [24], we have

$$(6.16) \quad \Delta_n \sim_{\mathcal{K}} (2\pi i)^{\frac{n(n+1)}{2}} \cdot \mathcal{G}(\varepsilon_{\mathcal{K}, f})^{\lceil \frac{n+1}{2} \rceil}.$$

In [24] n is assumed even, but the same argument gives the above result. Since H and H' are both definite unitary groups, the archimedean local integral Z_∞ is an algebraic number, rational over the field of definition of (the finite-dimensional representation) $\pi_\infty \otimes \pi'_\infty$. The discussion leading to §4.1.6 of [24] thus yields:

Corollary 6.17. *Assume π , π' , and \mathcal{K} satisfy hypothesis 6.12. Then there exists a non-zero constant $c_{\pi_\infty, \pi'_\infty}$ such that*

$$(6.18) \quad \Lambda(\pi, \pi') := c_{\pi_\infty, \pi'_\infty} \mathcal{G}(\varepsilon_{\mathcal{K}, f})^{\lceil \frac{n+1}{2} \rceil} (2\pi i)^{\frac{n(n+1)}{2}} \cdot \frac{L(\frac{1}{2}, \Pi_f \times \Pi'_f)}{L(1, \pi_f, Ad)L(1, \pi'_f, Ad)} \in \bar{\mathbb{Q}}$$

- For all $\sigma \in \text{Aut}(\mathbb{C})$,

$$(6.19) \quad \sigma(\Lambda(\pi, \pi')) = \Lambda(\sigma\pi, \sigma\pi')$$

The only difference with (4.1.6.2) of [24] is the inclusion of the factor $c_{\pi_\infty, \pi'_\infty}$ – since the Ichino-Ikeda-Neal Harris conjecture is only known up to a factor $c(\pi_\infty, \pi'_\infty)$ – and of the power of the Gauß sum is not omitted.

Theorem 1.3 gives a different expression for the numerator of (6.18). Before we compare the two expressions, we need the following non-vanishing result, which is essentially due to Wei Zhang.

Proposition 6.20. *Let π be an automorphic representation of $H(\mathbb{A})$ satisfying hypothesis 6.12. Then there exists an automorphic representation π' of $H'(\mathbb{A})$ satisfying hypothesis 6.12 and $\phi \in \pi$, $\phi' \in \pi'$, such that $P(\phi, \phi') \neq 0$ and $L(\frac{1}{2}, BC(\pi) \times BC(\pi')) \neq 0$.*

Proof. This is essentially Lemma 2.15 of [52]. Let p be a split place such that π_p is supercuspidal. Since the center of H is anisotropic, the central character of π_p is unitary. Let μ_p be a supercuspidal representation of $H'(\mathbb{Q}_p)$ with unitary central character. Since $H(\mathbb{Q}_p) \cong GL_n(\mathbb{Q}_p)$ and $H'(\mathbb{Q}_p) \cong GL_{n-1}(\mathbb{Q}_p)$, it is known that $\text{Hom}_{H'(\mathbb{Q}_p)}((\Pi_p \otimes \mu_p), \mathbb{C}) \neq 0$.

Since both π_p and μ_p are tempered, it follows from a result of Sakellaridis and Venkatesh (quoted as Theorem A.1 in [27]) that the local integral

$$\int_{H'(\mathbb{Q}_p)} f(h')f'(h')dh' \neq 0$$

for some matrix coefficients f, f' of π_p and μ_p , respectively. It follows that (the dual of) μ_p is weakly contained in the restriction to $H'(\mathbb{Q}_p)$ of π_p .

Now we assume μ_p is induced from an irreducible representation of $GL_{n-1}(\mathbb{Z}_p) \cdot Z$, where Z is the center of $H'(\mathbb{Q}_p)$; by the theory of types, there are supercuspidal representations with this property. Choose an irreducible component μ_∞ of the restriction to $H'(\mathbb{R})$ of (the finite-dimensional representation) π_∞ . Then we can apply Lemma 2.15 of [52] to obtain an automorphic representation π' of H' , whose archimedean component is μ_∞ and whose p -component is supercuspidal (a twist of μ_p), for which there exist $\phi \in \pi$ and $\phi' \in \pi'$ such that $P(\phi, \phi') \neq 0$. This implies the non-vanishing of the L -value by the main result of [53], provided we know that π' is unramified at all non-split places. \square

Lemma 6.21. *Let π be an irreducible representation of $H(\mathbb{A})$ when $n = 1$; in other words, π is a character of $U(1)(\mathbb{A})$. Then $L(s, \pi, Ad) = L(s, \varepsilon_{\mathcal{K}})$. In particular, $L(1, \pi_f, Ad) \sim_{\mathcal{K}} (2\pi i)\mathcal{G}(\varepsilon_{\mathcal{K}, f})$.*

Proof. This is an obvious calculation: The Lie algebra of H is one-dimensional and the adjoint representation of ${}^L G$ is trivial on the Langlands dual group \mathbb{C}^\times and is given by the quadratic character $\varepsilon_{\mathcal{K}}$. \square

Theorem 6.22. *Let π be an irreducible unitary cuspidal automorphic representation of $H(\mathbb{A})$ which is cohomological with respect to E_μ^{unt} . Assume π satisfies hypothesis 6.12. Then there are a non-zero complex constant $a(\pi_\infty)$, depending only on π_∞ , and an integer $g(n) \in \mathbb{Z}$, such that*

$$p(BC(\pi)) \sim_{\mathcal{K}\mathbb{Q}(\pi)} a(\pi_\infty)\mathcal{G}(\varepsilon_{\mathcal{K}, f})^{g(n)}L(1, \pi_f, Ad).$$

Proof. We prove this by induction on n . When $n = 1$ the Whittaker period $p(BC(\pi))$ is rational and the claim follows from 6.21.

Suppose the theorem is known for $n - 1$. Choose π and let π' be an automorphic representation of $H'(\mathbb{A})$ satisfying 6.20 relative to π . As π and π' both satisfy hypothesis 6.12, $\Pi = BC(\pi)$ and $\Pi' = BC(\pi')$ are cuspidal automorphic representations matching the conditions of Thm. 1.3. In particular, $p(BC(\pi))$ and $p(BC(\pi'))$ exist. By the very construction of π' , Lem. 2.1 and Lem. 4.7, $s_0 = \frac{1}{2}$ is critical for $L(s, BC(\pi) \times BC(\pi'))$, so $p(BC(\pi)_\infty, BC(\pi')_\infty)$ is well-defined. Now, let

$$\Lambda_W(\pi, \pi') := c_{\pi_\infty, \pi'_\infty} \mathcal{G}(\varepsilon_{\mathcal{K}, f})^{\lceil \frac{n+1}{2} \rceil} (2\pi i)^{\frac{n(n+1)}{2}} \cdot \frac{p(BC(\pi))p(BC(\pi'))p(BC(\pi)_\infty, BC(\pi')_\infty)}{L(1, \pi_f, Ad)L(1, \pi'_f, Ad)}.$$

We apply (6.18) and Thm. 1.3 to the pair (π, π') and conclude that $\Lambda_W(\pi, \pi') \in \bar{\mathbb{Q}}$. By [24], Prop. 2.6, $\sigma\Pi$ (resp. $\sigma\Pi'$) descends to $\sigma\pi$ (resp. $\sigma\pi'$), which are cuspidal automorphic representations satisfying hypothesis 6.12. Hence, using Prop. 6.20, (6.19) and Thm. 1.3 again,

$$(6.23) \quad \sigma(\Lambda_W(\pi, \pi')) = \Lambda_W(\sigma\pi, \sigma\pi'),$$

for all $\sigma \in \text{Aut}(\mathbb{C})$. But by induction, we can rewrite

$$\Lambda_W(\pi, \pi') \sim_{\mathcal{K}\mathbb{Q}(\pi')} \frac{p(BC(\pi))}{L(1, \pi_f, Ad)} \cdot [a(\pi'_\infty) c_{\pi_\infty, \pi'_\infty} \mathcal{G}(\varepsilon_{\mathcal{K}, f})^{\lceil \frac{n+1}{2} \rceil + g(n-1)} (2\pi i)^{\frac{n(n+1)}{2}} \cdot p(BC(\pi)_\infty, BC(\pi')_\infty)].$$

It follows from (6.23) that

$$p(BC(\pi)) \sim_{\mathcal{K}\mathbb{Q}(\pi)\mathbb{Q}(\pi')} L(1, \pi_f, Ad) \cdot [a(\pi'_\infty) c_{\pi_\infty, \pi'_\infty} \mathcal{G}(\varepsilon_{\mathcal{K}, f})^{\lceil \frac{n+1}{2} \rceil + g(n-1)} (2\pi i)^{\frac{n(n+1)}{2}} \cdot p(BC(\pi)_\infty, BC(\pi')_\infty)].$$

The term in brackets on the right-hand side depends only on π_∞ (since π'_∞ is an irreducible component of the restriction to $H'(\mathbb{R})$ of π_∞) and the degree n , whereas the left-hand side depends only on π , so we conclude by induction. \square

Remark 6.24. For a different approach to a similar (in fact more general) result, we refer to our forthcoming work with E. Lapid, [15].

Corollary 6.25. *Let Π and Π' be cuspidal automorphic representations of $G(\mathbb{A}_{\mathcal{K}})$ and $G'(\mathbb{A}_{\mathcal{K}})$ which are cohomological with respect to E_μ and E_λ , respectively. We assume $\Pi = BC(\pi)$ and $\Pi' = BC(\pi')$ where π and π' are irreducible unitary cuspidal automorphic representations of the definite unitary groups $H(\mathbb{A})$ and $H'(\mathbb{A})$, respectively. Suppose moreover that both π and π' satisfy hypothesis 6.12 and that the coefficients E_μ and E_λ satisfy the equivalent conditions of Lem. 2.1. Then for every critical point $s_0 = \frac{1}{2} + m$ of $L(s, \Pi \times \Pi')$ with $m \geq 0$, there are a non-zero complex constant $a(m, \Pi_\infty, \Pi'_\infty)$, depending only on the archimedean components of Π and Π' , and an integer $a(n)$, depending only on n , such that,*

$$L(\frac{1}{2} + m, \Pi_f \times \Pi'_f) \sim_{\mathcal{K}\mathbb{Q}(\pi)\mathbb{Q}(\pi')} a(m, \Pi_\infty, \Pi'_\infty) \mathcal{G}(\varepsilon_{\mathcal{K}, f})^{a(n)} L(1, \pi_f, Ad) L(1, \pi'_f, Ad).$$

Equivalently, for every critical point $s_0 = n - 1 + m$ of $L(s, R(M(\Pi) \otimes M(\Pi')))$ with $m \geq 0$,

$$L(n-1+m, R(M(\Pi) \otimes M(\Pi'))) \sim_{\mathcal{K}\mathbb{Q}(\pi)\mathbb{Q}(\pi')} a(m, \Pi_\infty, \Pi'_\infty) \mathcal{G}(\varepsilon_{\mathcal{K}, f})^{a(n)} L(1, \pi_f, Ad) L(1, \pi'_f, Ad).$$

Proof. Indeed, it follows from the Theorems 6.22 and 1.3 and the fact that $\mathbb{Q}(\Pi_f) \subseteq \mathbb{Q}(\pi)$ (resp. $\mathbb{Q}(\Pi'_f) \subseteq \mathbb{Q}(\pi')$) that we can define

$$a(m, \Pi_\infty, \Pi'_\infty) := a(\pi_\infty) a(\pi'_\infty) p(m, \Pi_\infty, \Pi'_\infty) \in \mathbb{C}^\times$$

and $a(n) := g(n) + g(n-1) \in \mathbb{Z}$. \square

6.6. Generalizations. The methods of this paper apply to pairs Π, Π' of representations when the critical values of their L -functions can be related directly to cup products of the cohomology classes they define. This is only possible when the coefficients satisfy the inequalities of Lemma 2.1, a condition that is equivalent to relations on the Hodge types of the corresponding motives that are summarized as 4.6.

We have seen in 4.12 that the Deligne periods of tensor products of motives satisfying these hypotheses can be expressed, up to a certain power of $2\pi i$, as products of terms denoted $P_{\leq s}(M)$ and $P_{\leq s'}(M')$, each occurring to the first power (for $1 \leq s \leq n-1$ and $1 \leq s' \leq n-2$). Deligne periods of tensor products are calculated more generally in §1.4 of [24]. As explained in [24], the motivic periods that occur in this calculation belong to a *tableau*, and it follows without difficulty that in all cases these Deligne periods are products of certain integral powers of the same $P_{\leq s}(M)$ and $P_{\leq s'}(M')$; the powers depend on the relative positions of the Hodge types of M and M' . In Optimistic Comparison 4.32 we

argue that $P_{\leq s}(M)$ and $P_{\leq s'}(M')$ can be identified with certain Petersson (square) norms of normalized holomorphic automorphic forms $P^{(i)}(\Pi)$ and $P^{(j)}(\Pi')$. Thus we would expect that the critical values of $L(s, \Pi \times \Pi')$ can always be expressed, up to algebraic factors, as powers of $2\pi i$ multiplied by powers of these Petersson inner products.

In fact, one can always find integers $N > n$ and Hecke characters $\chi_i, i = 1, \dots, N - n$; $\chi'_j, j = 1, \dots, N - n$, all obtained by base change from $U(1)$, such that the (tempered) Eisenstein representations

$$\Sigma = \Pi \boxplus \chi_1 \boxplus \chi_2 \boxplus \dots \boxplus \chi_{N-n}; \quad \Sigma' = \Pi' \boxplus \chi'_1 \boxplus \chi'_2 \boxplus \dots \boxplus \chi'_{N-n}$$

are cohomological and have coefficients satisfying the inequalities of Lemma 2.1. (In alluding to "base change" we are ignoring the parity issue that we have already seen in 2.5; all the parameters of the Eisenstein representations have to be either integers or half-integers, depending on the parity of N . But this can always be arranged.) Suppose for the moment that the assertion of Theorem 3.9 was valid for the pair (Σ, Σ') . Then we would find that, up to algebraic factors, the critical values of $L(s, \Sigma \times \Sigma')$ were powers of $2\pi i$ multiplied by

$$\prod_{r=1}^{N-1} P^{(r)}(\Sigma) \cdot \prod_{r'=1}^{N-2} P^{(r')}(\Sigma').$$

On the other hand, the L -function admits a factorization

$$L(s, \Sigma \times \Sigma') = L(s, \Pi \times \Pi') \cdot \prod_{i=1}^{N-n} L(s, \Pi' \otimes \chi_i) \cdot \prod_{j=1}^{N-n} L(s, \Pi \otimes \chi'_j) \cdot \prod_{i,j=1}^{N-n} L(s, \chi_i \cdot \chi_j).$$

Now we have expressions of the last three factors on the right-hand side in terms of $P^{(i)}(\Pi)$ and $P^{(j)}(\Pi')$ and CM periods of χ_i and χ'_j , and we can expect that $P^{(r)}(\Sigma)$ and $P^{(r')}(\Sigma')$ can also be expressed in terms of the periods of Π , Π' , and the auxiliary Hecke characters. In this way one would obtain an expression of the critical values of interest, namely those of the remaining term $L(s, \Pi \times \Pi')$, as powers of $P^{(i)}(\Pi)$ and $P^{(j)}(\Pi')$, as expected.

The problem is that the integral representation used in Theorem 3.9 does not converge when the automorphic form on GL_N is an Eisenstein series. In [26], Ichino and Yamana obtain the Rankin-Selberg product for Eisenstein representations of $GL_N \times GL_{N-1}$ as a regularized period integral. It seems to be difficult to interpret this regularized integral as a regularized cup product in rational cohomology.

Alternatively, assuming both Σ and Σ' descend to definite unitary groups, which must certainly be possible for appropriate choices of χ_i and χ'_j , we could apply the method described in Section 6.5 to obtain an expression for the central critical value $L(\frac{1}{2}, \Pi \times \Pi')$ analogous to that in Corollary 6.25, provided the χ_i and χ'_j could be chosen, with the given infinity types, so that the central critical values $L(\frac{1}{2}, \Pi' \otimes \chi_i)$ and $L(\frac{1}{2}, \Pi \otimes \chi'_j)$ were all non-zero. It would then be possible to deduce the expected expressions for more general central critical values from the expected generalization of recent results of Harder and Raghuram [19]. Unfortunately, proving the existence of such χ_i and χ'_j seems to be an extremely difficult problem.

Yet another method would be to ignore the cup product altogether. The Rankin-Selberg integral

$$(6.26) \quad \phi \otimes \phi' \mapsto I_m(\phi, \phi') := \int_{G'(\mathcal{K}) \backslash G'(\mathbb{A}_{\mathcal{K}})} \phi(g') \phi'(g') \|\det(g)\|^m dg$$

defines a $(\mathfrak{g}, K, G(\mathbb{A}_f))$ -invariant bilinear pairing on the space $\Pi_{(K)} \otimes \Pi'_{(K')}$ of $K \times K'$ -finite elements in $\Pi \otimes \Pi'$, indexed by the integer m (where this is defined by analytic continuation of the integral where necessary). On the other hand, its realization in cohomology determines a natural $\mathbb{Q}(\Pi_f)\mathbb{Q}(\Pi'_f)$ -rational structure on $\Pi_f \otimes \Pi'_f$, and therefore $L(\Pi, \Pi') := \text{Hom}_{(\mathfrak{g}, K, G(\mathbb{A}_f))}(\Pi_{(K)} \otimes \Pi'_{(K')}, \mathbb{C})$ also has a natural $\mathbb{Q}(\Pi_f)\mathbb{Q}(\Pi'_f)$ -rational structure. Under the so-called automatic continuity hypothesis, any element of $L(\Pi, \Pi')$ extends continuously to the smooth Fréchet completions of moderate growth. In this case, it is known that $L(\Pi, \Pi')$ is of dimension 1. Thus, there is a Rankin-Selberg period invariant $P(\Pi, \Pi') \in \mathbb{C}^\times$ such that $P(\Pi, \Pi')^{-1}I_m$ is $\mathbb{Q}(\Pi_f)\mathbb{Q}(\Pi'_f)$ -rational; moreover, these invariants can be defined consistently for all $\text{Aut}(\mathbb{C})$ -conjugates of Π, Π' . Verification of Deligne's conjecture then comes down to identifying these $P(\Pi, \Pi')$, whether or not the pairing I_m can be related to a cohomological cup product. If Π' is abelian automorphic, the $P(\Pi, \Pi')$ can be related as above to Petersson norms of holomorphic forms on unitary groups and periods of Hecke characters. It would seem that $P(\Pi, \Pi')$ factors in general as a product of a period invariant attached to Π and one attached to Π' , each independent of the other factor, but we see no way to prove this in general.

In a forthcoming joint paper with S. Yamana, we avoid these difficulties by applying a different combinatorial argument. We anticipate that our method will give a version of Theorem 6.10 without assuming the inequalities of Lemma 2.1.

REFERENCES

- [1] A. Ash, Non-square-integrable cohomology classes of arithmetic groups, *Duke J. Math.* **47** (1980) pp. 435–449
- [2] R. Beuzart-Plessis, *La conjecture locale de Gross-Prasad pour les représentations tempérées des groupes unitaires*, preprint (2012) arXiv:1205.2987v2.
- [3] A. Borel, J.-P. Serre, Corners and arithmetic groups, *Comm. Math. Helvet.* **48** (1973) pp. 436–491
- [4] A. Borel, N. Wallach, *Continuous cohomology, discrete subgroups and representations of reductive groups*, Ann. of Math. Studies, (Princeton Univ. Press, New Jersey, 2000)
- [5] A. Caraiani, Local-global compatibility and the action of monodromy on nearby cycles, *Duke Math. J.*, **161** (2012), 2311–2413.
- [6] L. Clozel, Motifs et Formes Automorphes: Applications du Principe de Fonctorialité, in *Automorphic forms, Shimura varieties, and L-functions*, Vol. I, Perspect. Math., vol. 10, eds. L. Clozel and J. S. Milne, (Ann Arbor, MI, 1988) Academic Press, Boston, MA, 1990, pp. 77–159.
- [7] L. Clozel, M. Harris, J.-P. Labesse, *Construction of automorphic Galois representations, I*, in L. Clozel et al., eds., *The stable trace formula, Shimura varieties, and arithmetic applications. Volume I: Stabilization of the trace formula* Boston: International Press (2011) pp. 497–527.
- [8] J. W. Cogdell, I. I. Piatetski-Shapiro, *Remarks on Rankin-Selberg convolutions*, in: *Contributions to Automorphic Forms, Geometry, and Number Theory*, eds. H. Hida, D. Ramakrishnan, F. Shahidi, Johns Hopkins University Press, Baltimore, 2004, pp. 255–278
- [9] P. Deligne, *Valeurs de fonctions L et périodes d'intégrales*, *With an appendix by N. Koblitz and A. Ogus* in: Proc. Sympos. Pure Math., Vol. XXXIII, part II, AMS, Providence, R.I., (1979), pp. 313–346
- [10] J. Franke, Harmonic analysis in weighted L_2 -spaces, *Ann. Sci. de l'ENS 4^e série tome 31* **2** (1998) pp. 181–279
- [11] J. Franke, J. Schwermer, A decomposition of spaces of automorphic forms, and the Eisenstein cohomology of arithmetic groups, *Math. Ann.*, **311** (1998), pp. 765–790
- [12] R. Goodman, N. Wallach, *Symmetry, representations, and invariants*, Graduate Texts in Mathematics, **255** (Springer, Dordrecht, 2009)
- [13] J. Gordon, Transfer to characteristic zero: Appendix to [51], *Duke Math J.* **156** (2011) pp. 220–227.

- [14] H. Grobner, Residues of Eisenstein series and the automorphic cohomology of reductive groups, *Compos. Math.* **149** (2013) pp. 1061–1090
- [15] H. Grobner, M. Harris, E. Lapid, Whittaker rational structures and special values of the Asai L -function, to appear in *Contemp. Math.* (2015)
- [16] H. Grobner, A. Raghuram, On some arithmetic properties of automorphic forms of GL_m over a division algebra, *Int. J. Number Theory* **10** (2014) pp. 963–1013
- [17] H. Grobner, A. Raghuram, On the arithmetic of Shalika models and the critical values of L -functions for GL_{2n} , with an appendix by Wee Teck Gan, *Amer. J. Math.* **136** (2014) pp. 675–728
- [18] G. Harder, *Some results on the Eisenstein cohomology of arithmetic subgroups of GL_n* , in: *Cohomology of Arithmetic Groups and Automorphic Forms*, eds. J.-P. Labesse, J. Schwermer, Lecture Notes in Math. **1447** (Springer, 1990) pp. 85–153
- [19] G. Harder, A. Raghuram, Eisenstein cohomology for GL_N and ratios of critical values of Rankin-Selberg L -functions - I, manuscript (2014)
- [20] R. N. Harris, The Refined Gross-Prasad Conjecture for Unitary Groups, *Int. Math. Res. Not.* (2012) doi: 10.1093/imrn/rns219
- [21] M. Harris, L -functions and periods of polarized regular motives, *J. Reine Angew. Math.* **483** (1997) pp. 75–161
- [22] M. Harris, *A simple proof of rationality of Siegel-Weil Eisenstein series*, in: *Eisenstein series and applications*, W.T. Gan, S.S. Kudla, Y. Tschinkel eds., *Progr. Math.* **258**, Birkhäuser Boston, Boston, 2008, pp. 149–186
- [23] M. Harris, *Cohomological automorphic forms on unitary groups, II: period relations and values of L -functions*, in J.-S. Li et al, eds. *Harmonic Analysis, Group Representations, Automorphic Forms and Invariant Theory*, Vol. **12**, Lecture Notes Series, Institute of Mathematical Sciences, National University of Singapore (2007) pp. 89–150
- [24] M. Harris, L -functions and periods of adjoint motives, *Alg. & Number Theory* **7** (2013) 117–155
- [25] M. Harris, J.-P. Labesse, Conditional base change for unitary groups, *Asian J. Math.* **8** (2004) 653–683
- [26] A. Ichino, S. Yamana, Periods of automorphic forms: the case of $(GL_{n+1} \times GL_n, GL_n)$, to appear in *Compos. Math.* (2015).
- [27] A. Ichino, W. Zhang, Spherical characters for a strongly tempered pair, appendix to [52]
- [28] H. Jacquet, I. I. Piatetski-Shapiro, J. Shalika, Conducteur des représentations du groupe linéaire, *Math. Ann.* **256** (1981) pp. 199–214
- [29] H. Jacquet, I. I. Piatetski-Shapiro, J. Shalika, Rankin-Selberg convolutions, *Amer. J. Math.* **105** (1983) pp. 367–464
- [30] H. Jacquet, J. Shalika, The Whittaker models of induced representations, *Pacific J. Math.* **109** (1983) pp. 107–120
- [31] T. Kaletha, A. Minguez, S. W. Shin, P.-J. White, *Endoscopic Classification of Representations: Inner Forms of Unitary Groups*, preprint arXiv: 1409.3731 (2014).
- [32] A. W. Knap, *Local Langlands Correspondence: The Archimedean Case*, in: Proc. Sympos. Pure Math., Vol. 55, part II, eds. U. Jannsen, S. L. Kleinman, J.-P. Serre, AMS, Providence R.I., 1994, pp. 393–410
- [33] J.-P. Labesse, Changement de base CM et séries discrètes, in: *On the Stabilization of the Trace Formula*, Vol. I, eds. L. Clozel, M. Harris, J.-P. Labesse, B.-C. Ngô, International Press, Boston, MA, 2011, pp. 429–470
- [34] J.-S. Li, J. Schwermer, On the Eisenstein cohomology of arithmetic groups, *Duke Math. J.* **123** (2004) pp. 141–169
- [35] J. Lin, Period relations for automorphic induction and applications, I, *Comptes Rendus Math.* **353** (2015) pp. 95–100
- [36] J. Mahnkopf, Cohomology of arithmetic groups, parabolic subgroups and the special values of automorphic L -Functions on $GL(n)$, *Journal de l’Institut de Mathématiques de Jussieu* **4** (2005) pp. 553–637
- [37] J. Mahnkopf, Modular symbols and values of L -functions on GL_3 , *J. reine angew. Math.* **497** (1998) pp. 91–112
- [38] C. Mœglin, J.-L. Waldspurger, *Spectral decomposition and Eisenstein series*, Cambridge Univ. Press (1995)
- [39] Ch. P. Mok, Endoscopic classification of representations of quasi-split unitary groups, *Memoires of the AMS* **235** (2014)

- [40] A. Raghuram, *Critical values of Rankin-Selberg L -functions for $GL_n \times GL_{n-1}$ and the symmetric cube L -functions for GL_2 . With an appendix by Chandrasheel Bhagwat*, preprint (2014)
- [41] A. Raghuram, On the Special Values of Certain Rankin-Selberg L -Functions and Applications to Odd Symmetric Power L -Functions of Modular Forms, *Int. Math. Res. Not.* Vol. 2010 (2010) pp. 334 – 372
- [42] A. Raghuram, F. Shahidi, On certain period relations for cusp forms on GL_n , *Int. Math. Res. Not.* Vol. 2008 (2008), doi:10.1093/imrn/rnn077.
- [43] J. Rohlfs, Projective limits of locally symmetric spaces and cohomology, *J. reine angew. Math.* **479** (1996) pp. 149–182
- [44] F. Shahidi, *Eisenstein series and Automorphic L -functions*, Colloquium publications **58**, AMS, (2010)
- [45] J. Schwermer, *Kohomologie arithmetisch definierter Gruppen und Eisensteinreihen*, Lecture Notes in Mathematics, **988**, Springer-Verlag (1983)
- [46] B. Sun, *The nonvanishing hypothesis at infinity for Rankin-Selberg convolutions*, preprint arXiv: 1307.5357 (2014)
- [47] E. Urban, Formes automorphes cuspidales pour $GL(2)$ sur un corps quadratique imaginaire. Valeurs spéciales de fonction L et congruences, *Compos. Math.* **99** (1995) pp. 283–324
- [48] J.-L. Waldspurger, Quelques propriétés arithmétiques de certaines formes automorphes sur $GL(2)$, *Comp. Math.* **54** (1985) pp. 121–171
- [49] H. Yoshida, On a conjecture of Shimura concerning periods of Hilbert modular forms, *Amer. J. Math.* **117** (1995) pp. 1019–1038
- [50] H. Yoshida, Motives and Siegel modular forms, *Amer. J. Math.* **123** (2001) pp. 1171–1197
- [51] Z. Yun, *Duke Math J.* **156** (2011) pp. 167–228
- [52] W. Zhang, Fourier transform and the global Gan-Gross-Prasad conjecture for unitary groups, *Ann. of Math.* **180** (2014) pp. 971–1049
- [53] W. Zhang, Automorphic period and the central value of Rankin-Selberg L -function, *JAMS* **27** (2014) pp. 541–612

HARALD GROBNER: FAKULTÄT FÜR MATHEMATIK, UNIVERSITÄT WIEN, OSKAR-MORGENSTERN-PLATZ 1, A-1090 WIEN, AUSTRIA
E-mail address: harald.grobner@univie.ac.at

MICHAEL HARRIS: UNIV PARIS DIDEROT, SORBONNE PARIS CITÉ, UMR 7586, INSTITUT DE MATHÉMATIQUES DE JUSSIEU-PARIS RIVE GAUCHE, CASE 247, 4 PLACE JUSSIEU F-75005, PARIS, FRANCE;; SORBONNE UNIVERSITÉS, UPMC UNIV PARIS 06, UMR 7586, IMJ-PRG, F-75005 PARIS, FRANCE;; CNRS, UMR7586, IMJ-PRG, F-75013 PARIS, FRANCE;; DEPARTMENT OF MATHEMATICS, COLUMBIA UNIVERSITY, NEW YORK, NY 10027, USA.